

Guarded Variable Automata over Infinite Alphabets

Walid Belkhir¹ and Yannick Chevalier² and Michael Rusinowitch¹

¹ INRIA Nancy–Grand Est & LORIA walid.belkhir@inria.fr, rusi@loria.fr

² Université Paul Sabatier & IRIT Toulouse ychevali@irit.fr

Abstract. We define *guarded variable automata* (GVAs), a simple extension of finite automata over infinite alphabets. In this model the transitions are labeled by letters or variables ranging over an infinite alphabet and guarded by conjunction of equalities and disequalities. GVAs are strictly more expressive than both finite memory automata (FMA) of Kaminski and Francez [9], and variable automata of Grumberg et al. [8]. They are closed under intersection, union, concatenation and Kleene operator, and their nonemptiness problem is decidable. We show that the simulation preorder of GVAs is decidable. Our proof relies on the characterization of the simulation by means of games and strategies.

1 Introduction

The simple and powerful formalism of finite automata is widely used for system specification and verification. Considerable efforts have been devoted to extend finite automata to infinite alphabets: finite memory automata [9], data automata [5], variable automata [8], fresh-variable automata [3], only to cite a few (see [10] for a survey). When developing formalisms over infinite alphabets, the main challenge is to preserve as much as possible useful properties such as compositionality (i.e. closure under basic operations) and decidability of basic problems such as nonemptiness, membership, universality, language containment, simulation, etc

The language containment problem is a particularly important one in applications like formal verification. For instance, whether an implementation is conform to a specification amounts to decide the containment $L(A) \subseteq L(B)$, where A (resp. B) is an automaton formalizing the behavior of the implementation (resp. specification), and $L(A)$ is the language of words recognized by A .

The containment problem for finite automata (FAs) can be solved by using determinization, in a complete but inefficient way. Moreover, for several classes of automata over infinite alphabets, the containment problem turned out to be undecidable. This is the case for finite memory

automata [11] and variable automata [8]. As a practical alternative approach, a simulation preorder can be employed to overapproximate the containment relation (e.g. [7]). Indeed, simulation-based techniques are sometimes more efficient. For instance a simulation between two finite automata can be computed in polynomial time. To our knowledge, simulation has not been studied for the classes of automata over infinite alphabets from [9] and [8].

Contributions. In this paper we define *guarded variable automata*, or GVAs, a natural extension of finite automata over infinite alphabets. In this model the transitions are labeled by letters or variables ranging over an infinite alphabets and guarded by conjunction of equalities and disequalities. Besides, some variables are refreshed in some states, that is, these variables can be released so that new letters can be bound to them. The potential applicability of our model in verification (e.g. model checking [4]) and service composition [1] follows from the fact that GVAs are closed under intersection, union, concatenation and Kleene operator. The nonemptiness problem is decidable for GVAs, and the membership is NP-Complete. However, their universality and containment problems are undecidable. We introduce a simulation preorder for GVAs and show its decidability. The proof relies on a game-theoretic characterization of simulation. Hence GVAs enjoy nice closure and decidability properties and we can show that they are more expressive than both finite memory automata and variable automata.

Related work. GVAs are closely related to the classes of automata in [9,8,3]. but these classes are strictly included in GVAs. Here we give a procedure to decide the simulation preorder for GVAs. Simulation has not been studied, to our knowledge, for finite memory automata and variable automata, among other models over infinite alphabets.

Paper organization. Sec. 2 recalls standard notions. Sec. 3 introduces the new class of guarded variable automata. Sec. 4 shows closure properties and decidability of nonemptiness for GVAs. Sec. 5 introduces the simulation preorder of GVAs and shows its decidability. Sec. 6 studies the expressiveness of GVAs with respect to FMAs and variable automata. Future work directions are given in Sec. 7. Missing proofs are provided in external appendices.

2 Preliminaries

Let $\mathcal{X}$ be a finite set of variables, Σ an infinite alphabet of letters. A substitution is an idempotent mapping $\{x_1 \mapsto \alpha_1, \dots, x_n \mapsto \alpha_n\} \cup$

$\bigcup_{a \in \Sigma} \{a \mapsto a\}$ with variables $x_1, \dots, x_n$ in $\mathcal{X}$ and $\alpha_1, \dots, \alpha_n$ in $\mathcal{X} \cup \Sigma$. We call $\{x_1, \dots, x_n\}$ its *proper domain*, and denote it by $\text{dom}(\sigma)$. We denote by $\text{Dom}(\sigma)$ the set $\text{dom}(\sigma) \cup \Sigma$. We denote by $\text{codom}(\sigma)$ the set $\{a \in \Sigma \mid \exists x \in \text{dom}(\sigma) \text{ s.t. } \sigma(x) = a\}$. If all the $\alpha_i, i = 1 \dots n$ are letters then we say that σ is ground. The empty substitution (*i.e.*, with an empty proper domain) is denoted by $\emptyset$. The set of substitutions from $\mathcal{X} \cup \Sigma$ to a set A is denoted by $\zeta_{\mathcal{X}, A}$, or by $\zeta_{\mathcal{X}}$, or simply by ζ if there is no ambiguity. If σ_1 and σ_2 are substitutions that coincide on the domain $\text{dom}(\sigma_1) \cap \text{dom}(\sigma_2)$, then $\sigma_1 \cup \sigma_2$ denotes their union in the usual sense. We define the function $\mathcal{V} : \Sigma \cup \mathcal{X} \longrightarrow \mathcal{P}(\mathcal{X})$ by $\mathcal{V}(\alpha) = \{\alpha\}$ if $\alpha \in \mathcal{X}$, and $\mathcal{V}(\alpha) = \emptyset$, otherwise. For a function $F : A \rightarrow B$, and $A' \subseteq A$, the restriction of F on A' is denoted by $F|_{A'}$. If $k \in \mathbb{N}$ then we let $[k] = \{1, \dots, k\}$.

A two-players game is a tuple $\langle \text{Pos}_E, \text{Pos}_A, M, p^* \rangle$, where $\text{Pos}_E, \text{Pos}_A$ are disjoint sets of positions: **Eloise's** positions and **Abelard's** positions. $M \subseteq (\text{Pos}_E \cup \text{Pos}_A) \times (\text{Pos}_E \cup \text{Pos}_A)$ is a set of *moves*, and p^* is the starting position. A strategy for the player **Eloise** is a function $\rho : \text{Pos}_E \rightarrow \text{Pos}_E \cup \text{Pos}_A$, such that $(\wp, \rho(\wp)) \in M$ for all $\wp \in \text{Pos}_E$. A (possibly infinite) play $\pi = \langle \wp_1, \wp_2, \dots \rangle$ follows a strategy ρ for player **Eloise** iff $\wp_{i+1} = \rho(\wp_i)$ for all $i \in \mathbb{N}$ such that $\wp_i \in \text{Pos}_E$. Let $\mathcal{W}$ be a (possibly infinite) set of plays. A strategy ρ is *winning* for **Eloise** from a set $S \subseteq \text{Pos}_E \cup \text{Pos}_A$ according to $\mathcal{W}$ iff every play starting from a position in S and following ρ belongs to $\mathcal{W}$.

3 Guarded variable automata

In this section we define formally the class of GVAs. It is an extension of FVAs [3] with logical constraints, called guards.

Definition 1. *The set $\mathbb{G}$ of guards over $\Sigma \cup \mathcal{X}$ is inductively defined as follows: $G := \text{true} \mid \alpha = \beta \mid \alpha \neq \beta \mid G \wedge G$, where $\alpha, \beta \in \Sigma \cup \mathcal{X}$. We write $\sigma \models g$ if a substitution σ satisfies a guard g .*

We notice that adding the disjunction operator to the guards would not increase the expressivity of our model, see Remark 1 in Appendix A.

A guard is atomic iff it is either **true**, an equality, or an inequality. Given a finite set $S = \{g_1, \dots, g_n\} \subset \mathbb{G}$ of atomic guards we denote by $\bigwedge S$ the guard $\bigwedge_{i=1}^n g_i$. In particular $\bigwedge \emptyset = \text{true}$. We let $\prod S = \{\bigwedge S' \mid \emptyset \subseteq S' \subseteq S\}$. The set of variables of the guard $G = \bigwedge S$, denoted by $\mathcal{V}(G)$, is defined by $\mathcal{V}(G) = \bigcup_{g \in S} \mathcal{V}(g)$ and $\mathcal{V}(x\theta y) = \mathcal{V}(x) \cup \mathcal{V}(y)$, $\theta \in \{=, \neq\}$. The application of a substitution σ to G is denoted by $\sigma(G)$. We shall write $\sigma \vdash G$ if there exists a substitution σ' such that $\sigma \models \sigma'(G)$.

Let $D \subseteq \mathcal{X}$ and $S \subseteq \mathbb{G}$ be a finite set of atomic guards, and let $G = \bigwedge S$. The *deletion* of the variables in D from G , denoted by G^D is defined by $G^D = \bigwedge \{g \in S \mid \mathcal{V}(g) \cap D = \emptyset\}$.

The formal definition of GVAs follows.

Definition 2. A GVA is a tuple $\mathcal{A} = \langle \Sigma, \mathcal{X}, Q, Q_0, \tau, \delta, F, \kappa \rangle$ where Σ is an infinite set of letters, $\mathcal{X}$ is a finite set of variables, Q is a finite set of states, $Q_0 \subseteq Q$ is a set of initial states, $\tau : \mathcal{X} \rightarrow \Sigma$ is a partial function called the initial assignment of variables, $\delta : Q \times (\Sigma_{\mathcal{A}} \cup \mathcal{X}) \times \mathbb{G} \rightarrow 2^Q$ is a transition function where $\Sigma_{\mathcal{A}}$ is a finite subset of Σ , $F \subseteq Q$ is a set of accepting states, and $\kappa : \mathcal{X} \rightarrow 2^Q$, called the refreshing function.

For a GVA $\mathcal{A}$, we shall denote by $\Sigma_{\mathcal{A}}$ (resp. $\mathbb{G}_{\mathcal{A}}$) the finite set of letters (resp. of atomic guards) that appear in the transition function of $\mathcal{A}$. For a path $\pi = q_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} q_n$ in $\mathcal{A}$, we define $\eta_0(\pi) = q_0$ and $\eta_f(\pi) = q_n$. Let $\pi' = q_n \xrightarrow{\alpha_{n+1}, g_{n+1}} \dots \xrightarrow{\alpha_m, g_m} q_m$ be another path in $\mathcal{A}$. The composition of the paths π and π' , denoted by $\pi \circ \pi'$, is the path $\pi'' = q_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} q_n \xrightarrow{\alpha_{n+1}, g_{n+1}} \dots \xrightarrow{\alpha_m, g_m} q_m$.

The semantics of guards has to be clarified: while taking a transition some variables in the guard of this transition may be free, i.e. not associated with letters, and hence they must be instantiated such that the guard holds. The formal definitions of configuration, run and recognized language follow.

Definition 3. Let $\mathcal{A} = \langle \Sigma, \mathcal{X}, Q, Q_0, \tau, \delta, F, \kappa \rangle$ be a GVA. A configuration is a pair (q, M) where $q \in Q$ and $M : \mathcal{X} \rightarrow \Sigma$ is a substitution. We define a transition relation over the configurations as follows: $(q_1, M_1) \xrightarrow{a} (q_2, M_2)$, where $a \in \Sigma$, iff there exist a substitution σ and a label $\alpha \in \Sigma \cup \mathcal{X}$ such that $q_2 \in \delta(q_1, \alpha, g)$ and $\text{dom}(\sigma) \cap \text{dom}(M_1) = \emptyset$ and either:

i) $\alpha \in \text{Dom}(M_1)$, $M_1(\alpha) = a$, $\text{dom}(\sigma) = \mathcal{V}(M_1(g))$, $(M_1 \uplus \sigma) \models g$ and $M_2 = (M_1 \uplus \sigma)|_D$, with $D = \text{Dom}(M_1 \uplus \sigma) \setminus \kappa^{-1}(q_2)$


or

ii) $\alpha \in (\mathcal{X} \setminus \text{Dom}(M_1))$, $\text{dom}(\sigma) = \{\alpha\} \cup \mathcal{V}(M_1(g))$, $(M_1 \uplus \sigma) \models g$ and $M_2 = (M_1 \uplus \sigma)|_D$, with $D = \text{Dom}(M_1 \uplus \sigma) \setminus \kappa^{-1}(q_2)$.

A finite word $w = w_1 w_2 \dots w_n \in \Sigma^*$ is recognized by $\mathcal{A}$ iff there exists a run $(q_0, M_0) \xrightarrow{w_1} (q_1, M_1) \xrightarrow{w_2} \dots \xrightarrow{w_n} (q_n, M_n)$, such that $M_0 = \tau$, $q_0 \in Q_0$ and $q_n \in F$. The set of words recognized by $\mathcal{A}$ is denoted by $L(\mathcal{A})$.

Example 1. Let $\mathcal{A}_1$ and $\mathcal{A}_2$ be the GVAs depicted on the right with κ_i the refreshing function of $\mathcal{A}_i$, $i = 1, 2$, and $\kappa_1(y) = \{p_0\}$ and $\kappa_2(x) = \{q_0\}$, and τ_i , $i = 1, 2$ their initial assignment with $\tau_1 = \tau_2 = \emptyset$. The language $L(\mathcal{A}_1)$ consists of all the words in Σ^* in which the last letter

is different than all the other letters. This language can be recognized by a variable automaton [8] but not by a FMA [9]. Notice that while making the first loop over p_0 , the variable x of the guard ($y \neq x$) must be instantiated. On the other hand, the language $L(\mathcal{A}_2) = \{w_1w_2 \cdots w_{2n} \mid w_i \in \Sigma, n \geq 1, \text{ and } w_{2i-1} \neq w_{2i}, \forall i \in [n]\}$ can be recognized by a FMA but not by a variable automaton.



4 Properties of guarded variable automata

We study the closure properties of GVAs and some basic decision problems. Firstly, we show that GVAs and GVAs with empty initial assignment recognize the same languages. The idea is to turn the initial assignment τ_0 into the guard $\phi = \bigwedge \{(z = a) \mid \tau_0(z) = a\}$ and propagate it in the transitions of the automaton, and remove from this guard the variables which are refreshed at each state.

Lemma 1. *For a GVA (with initial assignment) we can construct a GVA with empty initial assignment recognizing the same language.*

Theorem 1. *GVAs are closed under union, concatenation, Kleene operator and intersection. They are not closed under complementation.*

For the closure under union we simply take the disjoint union of the two GVAs. The closure under Kleene operator and concatenation follows from the fact that we can extend GVAs with ε -transitions (and trivial guards) and show that they have the same expressivity as GVAs. (Claim 4 in Appendix B.2). The closure under intersection is a consequence of the fact that computing the intersection of two GVAs amounts to computing their Cartesian product, which can be turned into a GVA (Appendix B.2). For the complementation, consider the language L_2 of all the words in which there is a letter that occurs at least twice. In fact, L_2 is GVA-recognizable [3]. The complement of L_2 consists of all the words in which all the letters are different, which is not GVA-recognizable since we need to compare each letter w_j to all the previous letters $w_i, i < j$. Thus we need an infinite number of states.

Despite GVAs are not closed under complementation, FAs can be complemented within the class of GVAs. That is, given a FA F there exists

a GVA $\mathcal{A}$ such that $L(\mathcal{A}) = \Sigma^* \setminus L(F)$, see Proposition 1 in Appendix B.2. It is worth mentioning that FAs cannot be complemented within the subclass of GVAs.

Decision procedures for GVAs. We study the decidability and complexity of classical decision problems: Nonemptiness (given $\mathcal{A}$, is $L(\mathcal{A}) \neq \emptyset$?), Membership (given a word w and $\mathcal{A}$, is $w \in L(\mathcal{A})$?), Universality (given $\mathcal{A}$, is $L(\mathcal{A}) = \Sigma^*$?), and Containment (given $\mathcal{A}_1$ and $\mathcal{A}_2$, is $L(\mathcal{A}_1) \subseteq L(\mathcal{A}_2)$?).

Theorem 2. *For GVAs, Membership is NP-complete, Universality and Containment are undecidable.*

The undecidability of Containment and Universality is a consequence of the undecidability of these problems for variable automata [8, Theorem 5]. However, the decidability of Containment if one of the GVAs is a finite automaton results from the fact that the intersection of the languages in this case is regular since the Cartesian product of a GVA and a FA yields a FA. The proof is the same as that of Lemma 17 of [3]. Hence,

Proposition 1. *The containment problems between a GVA and a FA are decidable.*

For Nonemptiness of GVAs (and more generally the reachability problems), we introduce first the so called *normal form* of GVAs. It is obtained by combining the *stretched* and *quotient* automaton defined below.

GVA's stretched form. It is useful to formalize the notion of a variable being *free* in a state.

Definition 4. *Let $\mathcal{A}$ be a GVA. A variable x is free in a state q of $\mathcal{A}$ if any run starting from the initial state and leading to q yields a configuration (q, M) in which $x \notin \text{Dom}(M)$.*

We define next the notion of *stretched form* for GVAs so that we can check whether a variable is free in a state without referring to the run.

Definition 5. *Let $\mathcal{A} = \langle \Sigma, \mathcal{X}, Q, Q_0, F, \delta, \kappa \rangle$ be a GVA. We define the stretched form of $\mathcal{A}$ to be the GVA $\mathcal{A}' = \langle \Sigma, \mathcal{X}, Q', Q'_0, F', \delta', \kappa' \rangle$ defined by:*

$$\begin{cases} Q' &= \{(q, X) \mid q \in Q \text{ and } X \subseteq \mathcal{X}\}, \\ Q'_0 &= \{(q, \mathcal{X}) \mid q \in Q_0\}, \\ F' &= \{(q, X) \mid q \in F \text{ and } X \subseteq \mathcal{X}\}. \end{cases}$$

The transition function δ' is defined by $(q', X') \in \delta'((q, X), \alpha, g)$, where $\alpha \in \Sigma \cup \mathcal{X}$ and $g \in \mathbb{G}$, if and only if, $X' = (X \setminus (\{\alpha\} \cup \mathcal{V}(g))) \cup \kappa^{-1}(q')$. Finally, the refreshing function κ' is defined by $\kappa'(x) = \{(q, X) \mid q \in \kappa(x)\}$.

Lemma 2. Let $\mathcal{A} = \langle \Sigma, \mathcal{X}, Q, Q_0, F, \delta, \kappa \rangle$ be a GVA and $\mathcal{A}' = \langle \Sigma, \mathcal{X}, Q', Q'_0, F', \delta', \kappa' \rangle$ be its stretched form. Then, $L(\mathcal{A}) = L(\mathcal{A}')$. Besides, a variable x is free in a state $(q, X) \in Q'$ iff $q \in X$.

Quotient GVA. Out of $\mathcal{A}$ we construct the GVA $\mathcal{A}_{/\approx}$ such that each state $\mathbf{q}$ of $\mathcal{A}_{/\approx}$ contains the guards which hold in $\mathbf{q}$ depending on the path leading from the starting state of $\mathcal{A}$ to $\mathbf{q}$.

Definition 6. Let $\mathcal{A} = \langle \Sigma, \mathcal{X}, Q, Q_0, F, \delta, \kappa \rangle$ be a GVA. The quotient automaton of $\mathcal{A}$ is the GVA $\mathcal{A}_{/\approx} = \langle \Sigma, \mathcal{X}, Q', Q'_0, F', \delta', \kappa' \rangle$ defined by:

$$\begin{cases} Q' &= \{(q, \phi) \mid q \in Q \text{ and } \phi \in \prod \mathbb{G}_{\mathcal{A}}\}, \\ Q'_0 &= \{(q, \text{true}) \mid q \in Q_0\}, \\ F' &= \{(q, \phi) \mid q \in F \text{ and } \phi \in \prod \mathbb{G}_{\mathcal{A}}\}, \end{cases}$$

The transition function δ' is defined by $(q', (g \wedge \phi)^{\kappa^{-1}(q')}) \in \delta'((q, \phi), \alpha, g)$ if, and only if, $q' \in \delta(q, \alpha, g)$, where $\alpha \in \mathcal{X} \cup \Sigma$ and $g \in \prod \mathbb{G}_{\mathcal{A}}$. The function κ' is defined by $\kappa'(x) = \{(q, \phi) \mid q \in \kappa(x) \text{ and } \phi \in \prod \mathbb{G}_{\mathcal{A}}\}$.

Lemma 3. There exists a run $q_0, M_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} q_n, M_n$ in $\mathcal{A}$ if, and only if, there exists a run $(q_0, \text{true}), M_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} (q_n, \phi_n), M_n$ in $\mathcal{A}_{/\approx}$ such that $M_i \models \phi_i$, for all $i = 0, \dots, n$.

For a GVA $\mathcal{A}$, the normal form of $\mathcal{A}$ is the GVA $\mathcal{A}'_{/\approx}$, where $\mathcal{A}'$ is the stretched form of $\mathcal{A}$.

Lemma 4. Let $\mathcal{A}$ be a GVA in the normal form. Let $\pi = (q_0, X_0, \phi_0) \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} (q_n, X_n, \phi_n)$ be a path in $\mathcal{A}$ with $\phi_0 = \text{true}$ and $q_0 \in Q_0$. Then, there is a run over π iff $\vdash \phi_n$ and $\vdash \phi_i \wedge g_{i+1}$, for all $i = 0, \dots, n-1$.

The following Corollary shows that to deal with the reachability in a GVA in the normal form, it suffices to consider simple paths.

Corollary 1. Let $\mathcal{A}$ be a GVA in the normal form. Let $\pi_1 \circ \pi_2 \circ \pi_3$ be a path in $\mathcal{A}$ with $\eta_f(\pi_1) = \eta_0(\pi_2) = \eta_f(\pi_2) = \eta_0(\pi_3)$. If there is an accepting run in $\mathcal{A}$ over $\pi_1 \circ \pi_2 \circ \pi_3$ then there is an accepting run over $\pi_1 \circ \pi_3$ as well.

Summing up the results shown so far, we are ready to prove the main result of this section.

Theorem 3. *Nonemptiness for GVAs is decidable.*

Proof. Let $\mathcal{A}$ be a GVA in stretched form. Let $\mathcal{A}_{/\approx}$ be its quotient automaton. From Lemma 3 it follows that $L(\mathcal{A}) = \emptyset$ iff $L(\mathcal{A}_{/\approx}) = \emptyset$. It follows from Corollary 1 that to check the nonemptiness of $\mathcal{A}_{/\approx}$ it is sufficient to look for an accepting run over a *simple* path of $\mathcal{A}_{/\approx}$. $\square$

5 Simulations for GVAs

We define and study the simulation preorder for GVAs, it is a generalization of the standard simulation for FAs. The idea is that the simulation for GVAs is over pairs of configurations. To simplify the presentation, we shall only consider in this section GVAs in which there is a unique initial state and all the states are accepting. The formal definition of simulation is given in Def. 11, Appendix C. In order to study the decidability of the simulation, we provide next an equivalent game-theoretic definition in which we make explicit the evolution of the configurations.

Definition 7. Let $\mathcal{A}_1 = \langle \Sigma, \mathcal{X}_1, Q_1, q_0^1, \tau_0^1, \delta_1, F_1, \kappa_1 \rangle$ and $\mathcal{A}_2 = \langle \Sigma, \mathcal{X}_2, Q_2, q_0^2, \tau_0^2, \delta_2, F_2, \kappa_2 \rangle$ be two GVAs where $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$. Let Pos be the set of positions reachable from $p^* = ((\tau_0^1, q_0^1), (\tau_0^2, q_0^2))_A$ by the set of moves $M = M_A \cup M_E$, where:

$$\begin{aligned} M_A = \{ & ((\sigma_1, q_1), \tau_2)_A \rightarrow (((\sigma_1 \uplus \gamma)|_D, q'_1), \tau_2, (\sigma_1 \uplus \gamma, \alpha))_E \\ & \mid q'_1 \in \delta_1(q_1, \alpha, g_1) \\ & \text{and } D = \text{Dom}(\sigma_1 \uplus \gamma) \setminus \kappa_1^{-1}(q'_1) \\ & \text{and } \sigma_1 \uplus \gamma \models g_1 \\ & \text{and } \gamma : \mathcal{V}(\sigma_1(\alpha)) \cup \mathcal{V}(\sigma_1(g_1)) \rightarrow \Sigma \} \\ M_E = \{ & ((\sigma_1, q_1), (\sigma_2, q_2), (\sigma_3, \alpha))_E \rightarrow ((\sigma_1, q_1), ((\sigma_2 \uplus \gamma_2)|_{D_2}, q'_2))_A \\ & \mid q'_2 \in \delta_2(q_2, \beta, g_2) \\ & \text{and } D_2 = \text{Dom}(\sigma_2 \uplus \gamma_2) \setminus \kappa_2^{-1}(q'_2) \\ & \text{and } \gamma_1(\sigma_3(\alpha)) = \gamma_2(\sigma_2(\beta)) \\ & \text{and } \gamma_2 \models \sigma_2(g_2) \\ & \text{and } \gamma_2 : \mathcal{V}(\sigma_2(\beta)) \cup \mathcal{V}(\sigma_2(g_2)) \rightarrow \Sigma \} \end{aligned}$$

We let $\text{Pos}_E = \text{Pos} \cap (\zeta_{\mathcal{X}_1} \times Q_1) \times (\zeta_{\mathcal{X}_2} \times Q_2) \times (\zeta_{\mathcal{X}_1} \times \Sigma \cup \mathcal{X})$ and $\text{Pos}_A = \text{Pos} \cap (\zeta_{\mathcal{X}_1} \times Q_1) \times (\zeta_{\mathcal{X}_2} \times Q_2)$. The simulation game of $\mathcal{A}_1$ by $\mathcal{A}_2$, denoted by $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$, is the two-players game $\langle \text{Pos}_E, \text{Pos}_A, M, p^* \rangle$. As usual, any infinite play is winning for *Eloise*, and any finite play is losing for the player who cannot move. And thus we write $\mathcal{A}_1 \trianglelefteq \mathcal{A}_2$.

The simulation problem for GVAs is the following: given two GVAs $\mathcal{A}_1$ and $\mathcal{A}_2$, is $\mathcal{A}_1 \trianglelefteq \mathcal{A}_2$?

5.1 Decidability of the simulation problem

In this section we show that the simulation problem is decidable. The idea is that this problem can be reduced to a simulation problem over the same GVAs in which the two players instantiate the variables from a *finite* set of letters, as proven in Proposition 2.

Definition 8. Let $\mathcal{A}_1 = \langle \Sigma, \mathcal{X}_1, Q_1, q_0^1, \tau_1, \delta_1, F_1, \kappa_1 \rangle$ and $\mathcal{A}_2 = \langle \Sigma, \mathcal{X}_2, Q_2, q_0^2, \tau_2, \delta_2, F_2, \kappa_2 \rangle$ be two GVAs. We define $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ to be the game obtained by restricting the codomain of γ to C_0 in the rules of *Eloise* M_E and *Abelard* M_A in Def. 7, where $C_0 = \Sigma_{\mathcal{A}_1} \cup \Sigma_{\mathcal{A}_2} \cup (\mathcal{X}_1 \times \mathcal{X}_2) \cup (\mathcal{X}_2 \times \mathcal{X}_1) \cup \text{codom}(\tau_1) \cup \text{codom}(\tau_2)$.

The following Lemma states an immediate property of the game $\overline{\mathcal{G}}$.

Lemma 5. Let $\mathcal{A}_1, \mathcal{A}_2$ be two GVAs. Then, the game $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ is finite.

In order to prove Proposition 2 we need to introduce the notion of coherence between substitutions and between game positions. The coherence relation was introduced in [3], we reproduce it here.

Definition 9. Let C be a finite subset of Σ . The coherence relation $\bowtie_C \subseteq \zeta \times \zeta$ between substitutions is defined by $\bar{\sigma} \bowtie_C \sigma$ iff the three following conditions hold:

1. $\text{dom}(\bar{\sigma}) = \text{dom}(\sigma)$,
2. If $\bar{\sigma}(x) \in C$ then $\bar{\sigma}(x) = \sigma(x)$, and if $\sigma(x) \in C$, then $\bar{\sigma}(x) = \sigma(x)$, for any variable $x \in \text{dom}(\sigma)$, and
3. for any variables $x, y \in \text{dom}(\sigma)$, $\bar{\sigma}(x) = \bar{\sigma}(y)$ iff $\sigma(x) = \sigma(y)$.

The definition of the coherence between game positions, still denoted by $\bowtie_C$, follows.

Definition 10. Let C be a finite subset of Σ , and $\mathcal{A}_1 = \langle \Sigma, \mathcal{X}_1, Q_1, q_0^1, \tau_1, \delta_1, F_1, \kappa_1 \rangle$, and $\mathcal{A}_2 = \langle \Sigma, \mathcal{X}_2, Q_2, q_0^2, \tau_2, \delta_2, F_2, \kappa_2 \rangle$ be two GVAs s.t. $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$. Let Pos_E (resp. Pos_A) be the set of *Eloise*'s (resp. *Abelard*'s) positions in the game $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$. Then we define the relation: $\bowtie_C \subseteq \text{Pos}_A \times \text{Pos}_A \cup \text{Pos}_E \times \text{Pos}_E$ by:

- For any $\sigma_i, \bar{\sigma}_i$ of proper domain included in $\mathcal{X}_i$ ($i = 1, 2$) we have:
 $((\bar{\sigma}_1, q_1), (\bar{\sigma}_2, q_2))_A \bowtie_C ((\sigma_1, q_1), (\sigma_2, q_2))_A$ iff $(\bar{\sigma}_1 \uplus \bar{\sigma}_2) \bowtie_C (\sigma_1 \uplus \sigma_2)$.
- For any $\sigma_i, \bar{\sigma}_i$ of proper domain included in $\mathcal{X}_i$ ($i = 1, 2$), for any substitutions $\sigma, \bar{\sigma}$ with proper domain included in $\mathcal{X}_1$, we have:
 $((\bar{\sigma}_1 \cup \bar{\sigma}) \uplus \bar{\sigma}_2) \bowtie_C ((\sigma_1 \cup \sigma) \uplus \sigma_2)$ iff
 $((\bar{\sigma}_1, q_1), (\bar{\sigma}_2, q_2), (\bar{\sigma}, \alpha))_E \bowtie_C ((\sigma_1, q_1), (\sigma_2, q_2), (\sigma, \alpha))_E$.

Now we are ready to show that the games $\mathcal{G}$ and $\overline{\mathcal{G}}$ are equivalent in the following sense:

Proposition 2. *Let $\mathcal{A}_1 = \langle \Sigma, \mathcal{X}_1, Q_1, q_0^1, \tau_1, \delta_1, F_1, \kappa_1 \rangle$ and $\mathcal{A}_2 = \langle \Sigma, \mathcal{X}_2, Q_2, q_0^2, \tau_2, \delta_2, F_2, \kappa_2 \rangle$ be two GVAs. Then, **Eloise** has a winning strategy in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ iff she has a winning strategy in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$.*

Proof. Up to renaming of variables, we can assume that $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$.

For the direction " $\Rightarrow$ " we show that out of a winning strategy of **Eloise** in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ we construct a winning strategy for her in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$. For this purpose, we show that each move of **Abelard** in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ can be mapped to an **Abelard** move in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$, and that **Eloise** response in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ can be actually mapped to an **Eloise** move in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$. This mapping defines a relation $\mathcal{R}$ between the positions of $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ and the positions of $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$. Formally,

$$\mathcal{R} \subseteq \text{Pos}_E(\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)) \times \text{Pos}_E(\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)) \cup \\ \text{Pos}_A(\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)) \times \text{Pos}_A(\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)),$$

such that if $(\bar{\wp}, \wp) \in \mathcal{R}$, and the move $\bar{\wp} \rightarrow \bar{\wp}'$ in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ is mapped to $\wp \rightarrow \wp'$ in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$, or $\wp \rightarrow \wp'$ in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ is mapped to $\bar{\wp} \rightarrow \bar{\wp}'$ in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$, then $(\bar{\wp}', \wp') \in \mathcal{R}$. Furthermore, we impose that the following invariant (Inv- $\bowtie$) holds: If $(\bar{\wp}, \wp) \in \mathcal{R}$ then $\bar{\wp} \bowtie_C \wp$, where $C = \Sigma_{\mathcal{A}_1} \cup \Sigma_{\mathcal{A}_2} \cup \text{codom}(\tau_1) \cup \text{codom}(\tau_2)$. We recall that the variables in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ are instantiated from the finite set of letters $C_0 = C \cup (\mathcal{X}_1 \times \mathcal{X}_2) \cup (\mathcal{X}_2 \times \mathcal{X}_1)$.

The main part of the proof consists in finding the right way to relate the instantiation of the variables in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ and $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$. For this purpose, given two sets S_1 and S_2 of alphabets such that $S_1 \cap S_2 = C \neq \emptyset$, we define a function $\Theta_C^{S_1, S_2} : \xi_{\mathcal{X}, S_1} \times \xi_{\mathcal{X}, S_1} \times \xi_{\mathcal{X}, S_2} \rightarrow \xi_{\mathcal{X}, S_2}$, that given three substitutions M, γ and M' such that $\text{dom}(M) \cap \text{dom}(\gamma) = \emptyset$ and $M \bowtie_C M'$, constructs a substitution $\gamma' = \Theta_C^{S_1, S_2}(M, \gamma, M')$ such that $M \uplus \gamma \bowtie_C M' \uplus \gamma'$.

To show how to use Θ , assume $\wp = ((\sigma_1, q_1), (\sigma_2, q_2))_A$ is a position in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$, and γ is an instantiation made by **Abelard** from $\wp$ (i.e. γ in the move M_A of Def. 7). Assume also that $\bar{\wp} = ((\bar{\sigma}_1, q_1), (\bar{\sigma}_2, q_2))_A$ is a position in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ such that $(\bar{\wp}, \wp) \in \mathcal{R}$. Then, **Abelard**'s instantiation $\bar{\gamma}$ from $\bar{\wp}$ is defined by $\bar{\gamma} = \Theta_C^{\Sigma, C_0}(\sigma_1 \uplus \sigma_2, \gamma, \bar{\sigma}_1 \uplus \bar{\sigma}_2)$.

For the other direction, **Eloise**'s instantiation of the variables in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ from Σ is related to **Eloise**'s instantiation of the variables in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ from C_0 by following the same principle. Following this construction, we ensure that the invariant (Inv- $\bowtie$) is always maintained.

The proof of the direction $(\Leftarrow)$ is similar to the one of $(\Rightarrow)$: we follow the same instantiation principle and we keep the same definition of the $\bowtie$ -coherence. $\square$

It follows from Lemma 5 and Proposition 2:

Theorem 4. *The simulation problem is decidable for GVAs.*

6 Comparison with related classes of automata

In this section we argue that GVAs are strictly more expressive than variable automata [8] and FMA [9]. The latter models are incomparable. Firstly, we recall that variable automata is an extension of FA in which the transitions are labeled by letters or variables among $\mathcal{X} \cup \{y\}$. Given a variable automaton $\mathcal{B}$, the main idea is that the variables in $\mathcal{X} \cup \{y\}$ range over $\Sigma \setminus \Sigma_{\mathcal{B}}$, where $\Sigma_{\mathcal{B}}$ is the finite set of the letters appearing in the transition function of $\mathcal{B}$. Besides, the instantiations of y differ from the instantiations of the variables in $\mathcal{X}$. Out of $\mathcal{B}$ we can construct a GVA $\mathcal{A}$ recognizing the same language, Lemma 8 in Appendix D.1. In fact, $\mathcal{A}$ can mimic $\mathcal{B}$ by turning the above constraints on the variables $\mathcal{X} \cup \{y\}$ into guards, and refreshing only the variable y in the source states of the transitions labeled with y . We emphasize that this construction does not imply directly that the simulation for variable automata is decidable although it is decidable for GVAs.

Secondly, we recall that a FMA is a 7-tuple $\mathcal{F} = \langle \Sigma, k, Q, q_0, \tau, \varrho, \delta, F \rangle$ where $k \in \mathbb{N}^+$ is the number of registers, Q is a finite set of states, $q_0 \in Q$ is the initial state, $\tau : [k] \rightarrow \Sigma$ is a partial function called the *initial assignment*, $\varrho : Q \rightarrow [k]$ is a partial function called the *reassignment*, $\delta : Q \times [k] \rightarrow \mathcal{P}(Q)$ is called the *transition function*, and $F \subseteq Q$ is the set of final states. Intuitively, given a current assignment $R : [k] \rightarrow \Sigma$, when a FMA $\mathcal{F}$ is in state q and reads a symbol a , then $\mathcal{F}$ changes its state into q' if $q' \in \delta(q, i)$ provided that $R(i)$ equals a . If $a \notin \text{codom}(R)$, then $\mathcal{F}$ rewrites a into the register $\varrho(p)$ and changes the state into q' if $q' \in \delta(q, \varrho(q))$. The formal definition of the run of FMA is in Appendix D.

We can show that for every FMA with k registers and n states, we can construct a GVA with $k+1$ variables and $O(n \cdot (k+1)!)$ states recognizing the same language, Lemma 9 in Appendix D.2. The idea is that k variables are used to mimic the k registers and the remaining variable is used to store the current letter. In fact, this correspondence between the k registers and the $k+1$ variables can change depending on the transitions of the FMA.

Finally, since the language $L(\mathcal{A}_1)$ (resp. $L(\mathcal{A}_2)$) of Example 1 can not be recognized by any FMA (resp. any variable automaton) then:

Theorem 5. *GVAs are strictly more expressive than FMA and variable automata.*

7 Conclusion

It is worth investigating the decidability of the containment problem for the subclass of GFVAs without disequalities in the guards. We also plan to apply our result on GFVAs simulation to the synthesis of web service composition. In this context, disequalities should be useful to express security policy enforcement on services [6,2].

References

1. L. Akroun, B. Benatallah, L. Nourine, and F. Toumani. On decidability of simulation in data-centric business protocols. In *Business Process Management Workshops*, volume 132, pages 352–363. Springer, 2013.
2. M. Bartoletti, P. Degano, G. L. Ferrari, and R. Zunino. Model checking usage policies. To appear in *Math. Stuct. Comp. Sci.*, abridged version in TGC 2008. LNCS.
3. W. Belkhir, Y. Chevalier, and M. Rusinowitch. Fresh-variable automata for service composition. <http://arxiv.org/abs/1302.4205>.
4. B. Berard, M. Bidoit, A. Finkel, F. Laroussinie, A. Petit, L. Petrucci, and P. Schnoebelen. *Systems and Software Verification: Model-Checking Techniques and Tools*. Springer Publishing Company, Incorporated, 1st edition, 2010.
5. M. Bojanczyk, A. Muscholl, T. Schwentick, L. Segoufin, and C. David. Two-variable logic on words with data. In *LICS*, pages 7–16. IEEE, 2006.
6. P. Degano, G. L. Ferrari, and G. Mezzetti. Nominal automata for resource usage control. In *CIAA*, pages 125–137, 2012.
7. D. L. Dill, A. J. Hu, and H. Wong-Toi. Checking for language inclusion using simulation preorders. In *CAV '91*, pages 255–265, London, 1992. Springer-Verlag.
8. O. Grumberg, O. Kupferman, and S. Sheinvald. Variable automata over infinite alphabets. In *LATA10*, volume 6031 of *LNCS*, pages 561–572. Springer, 2010.
9. M. Kaminski and N. Francez. Finite-memory automata. *Theor. Comput. Sci.*, 134(2):329–363, 1994.
10. A. Manuel and R. Ramanujam. Automata over infinite alphabets. *World Scientific Review*, 9:329–363, 2011.
11. F. Neven, T. Schwentick, and V. Vianu. Finite state machines for strings over infinite alphabets. *ACM Trans. Comput. Log.*, 5(3):403–435, 2004.

Appendices

A Remarks for Section 3

Remark 1. It is possible to consider the disjunction operation ($\vee$) in the guards but one can show that any GVA, in which the guards contain the disjunction, can be turned into a GVA recognizing the same language and in which the guards are without disjunctions, i.e. in the set $\mathbb{G}$ defined earlier. But we must be careful about the instantiation of the potential free variables present in the guard: a transition $q_1 \xrightarrow{\alpha, g_1 \vee g_2} q_2$ can be equivalently turned into two transitions $q_1 \xrightarrow{\alpha, g_1 \wedge E_1} q_2$ and $q_1 \xrightarrow{\alpha, g_2 \wedge E_2} q_2$, where $E_1 = \bigwedge_{x \in S_1} (x = x)$ (resp. $E_2 = \bigwedge_{y \in S_2} (y = y)$) and $S_1 = \mathcal{V}(g_2) \setminus \mathcal{V}(g_1)$ (resp. $S_2 = \mathcal{V}(g_1) \setminus \mathcal{V}(g_2)$).

B Proofs for Section 4

B.1 GVAs with empty initial assignment

Claim 1 *Let g be a guard, σ a substitution and $D \subseteq \mathcal{X}$. If $\sigma \models g$ then $\sigma|_{\bar{D}} \models g^D$ where $\bar{D} = \text{Dom}(\sigma) \setminus D$.*

Proof. Assume $g = \bigwedge_{i \in I} g_i$ where every g_i is an atomic guard. The proof is by induction on $|D|$. If $D = \emptyset$, then the proof is trivial since $\bigwedge \emptyset = \text{true}$. For the induction case, assume $D = \{x\} \uplus D'$. On the one hand, we have $\bar{D} = \text{Dom}(\sigma) \setminus D = \text{Dom}(\sigma) \setminus (\{x\} \uplus D') = (\text{Dom}(\sigma) \setminus \{x\}) \setminus D'$. On the other hand, $g^D = g^{\{x\} \uplus D'} = (g^{\{x\}})^{D'}$. If $\sigma \models g$, from the induction hypothesis we have $\sigma|_{\text{Dom}(\sigma) \setminus \{x\}} \models g^{\{x\}}$ and $\sigma|_{(\text{Dom}(\sigma) \setminus \{x\}) \setminus D'} \models (g^{\{x\}})^{D'}$. $\square$

Lemma 1. *For a GVA (with initial assignment) we can construct a GVA with empty initial assignment recognizing the same language.*

Proof. Let $\mathcal{A} = \langle \Sigma, \mathcal{X}, Q, q_0, \tau_0, \delta, F, \kappa \rangle$ be a GVA. Define

$$\phi = \bigwedge \{(z = a) \mid \tau_0(z) = a\} \quad (1)$$

Let $\xi = \xi_{\text{dom}(\tau_0), \text{codom}(\tau_0)}$. We let $\mathcal{A}' = \langle \Sigma, \mathcal{X}, Q', q'_0, F', \delta', \kappa' \rangle$ defined by:

$$\begin{cases} Q' = \{(q, \tau) \mid q \in Q \text{ and } \tau \in \xi\} \\ q'_0 = (q_0, (\tau_0)|_{\text{dom}(\tau_0) \setminus \kappa^{-1}(q_0)}) \\ F' = \{(q, \tau) \mid q \in F \text{ and } \tau \in \xi\} \end{cases}$$

The transition function δ' is defined by:

$$(q', \tau_{|D}) \in \delta'((q, \tau), \alpha, g \wedge \phi_{|dom(\tau)}) \quad \text{iff} \quad q' \in \delta(q, \alpha, g),$$

where $\alpha \in \Sigma \cup \mathcal{X}$ and $g \in \mathbb{G}$ and $\tau \in \xi$ and $D = dom(\tau) \setminus \kappa^{-1}(q')$. Finally,

$$\kappa'(x) = \{(q, \tau) \mid q \in \kappa(x) \text{ and } \tau \in \xi\}.$$

The following Claim is straightforward.

Claim 2 *We have that $\tau_0 \models \phi$.*

Claim 3 *Let $D_0 = dom(\tau_0) \setminus \kappa^{-1}(q_0)$ and let $w = w_1 \dots w_n$ be a word in Σ^n . There exists a run*

$$q_0, M_0 \xrightarrow{\alpha_1, g_1} q_1, M_1 \rightarrow \dots \xrightarrow{\alpha_n, g_n} q_n, M_n$$

in $\mathcal{A}$ over w , where $M_0 = (\tau_0)_{|D_0}$ if and only if, there exists a run

$$(q_0, \tau_0), M'_0 \xrightarrow{\alpha_1, g_1 \wedge \phi_1} (q_1, \tau_1), M'_1 \rightarrow \dots \xrightarrow{\alpha_n, g_n \wedge \phi_n} (q_n, \tau_n), M'_n$$

in $\mathcal{A}'$ over w , where $M'_0 = \emptyset$, $\tau_0 = \tau_0|_{D_0}$, and for all $i = 0, \dots, n-1$, $\phi_{i+1} \stackrel{def}{=} \phi_{|dom(\tau_i)}$, and the following three invariants hold:

$$\tau_i \models \phi_{i+1}, \tag{Inv-1}$$

$$\tau_{i+1} \subseteq M'_{i+1}, \tag{Inv-2}$$

$$M_{i+1} = M'_{i+1}, \tag{Inv-3}$$

Proof (Of Claim 3). The proof is by induction on $n \geq 1$, the length of the run, in both directions.

– Base case $n = 1$.

$\Rightarrow$) Let

$$\begin{aligned} M_1 &\stackrel{def}{=} (M_0 \uplus \sigma)_{|D}, \text{ where } dom(\sigma) = \mathcal{V}(M_0(g_1)) \cup \mathcal{V}(M_0(\alpha_1)) \\ &\text{and } D = Dom(M_0 \uplus \sigma) \setminus \kappa^{-1}(q_1) \\ &\text{and } M_0 \uplus \sigma \models g_1 \\ &\text{and } (M_0 \uplus \sigma)(\alpha_1) = w_1 \end{aligned}$$

We must show the existence of a substitution σ' such that:

$$\begin{aligned} M'_1 &= (M'_0 \uplus \sigma')|_{D'}, \text{ where} \\ \text{dom}(\sigma') &= \mathcal{V}(M'_0(g_1 \wedge \phi_1)) \cup \mathcal{V}(M'_0(\alpha_1)) \\ \text{and } D' &= \text{Dom}(M'_0 \uplus \sigma') \setminus \kappa^{-1}(q_1) \\ \text{and } M'_0 \uplus \sigma' &\models g_1 \wedge \phi_1 \\ \text{and } (M'_0 \uplus \sigma')(\alpha_1) &= w_1 \end{aligned}$$

Since $M'_0 = \emptyset$, then M'_1 can be rewritten as

$$\begin{aligned} M'_1 &= \sigma'|_{D'}, \text{ where } \text{dom}(\sigma') = \mathcal{V}(g_1 \wedge \phi_1) \cup \mathcal{V}(\alpha_1) \\ \text{and } D' &= \text{Dom}(\sigma') \setminus \kappa^{-1}(q_1) \\ \text{and } \sigma' &\models g_1 \wedge \phi_1 \\ \text{and } \sigma'(\alpha_1) &= w_1 \end{aligned}$$

We take $\sigma' \stackrel{\text{def}}{=} M_0 \uplus \sigma$, and we must show that $M_0 \uplus \sigma \models g_1 \wedge \phi_1$, that is, we must show $M_0 \uplus \sigma \models \phi_1$. From Claim 2 we have $\tau_0 \models \phi$, hence $\rho_0 = M_0 = (\tau_0)|_{D_0} \models \phi^{\kappa^{-1}(q_0)} = \phi_1$, $D = D'$ and $M_1 = M'_1$. Thus the invariants (Inv-1) and (Inv-3) hold. To prove (Inv-2), on the one hand we have $\tau_0 = M_0$ and $\rho_1 = (\tau_0)_{\text{Dom}(\tau) \setminus \kappa^{-1}(q_1)}$. On the other hand, $M'_1 = M_1 = (M_0 \uplus \sigma)|_{\text{Dom}(M_0 \uplus \sigma) \setminus \kappa^{-1}(q_1)}$. Therefore, $\tau_1 \subseteq M'_1$.

- $\Leftarrow$) Let M'_1 (as in the previous direction) and we find a substitution σ such that M_1 exists. We take $\sigma = \sigma' \setminus M_0$. The proof that the invariants hold is similar to the previous direction.
- Induction case. Assume the claim holds up to n . Let us prove the equivalence for $n + 1$.
- $\Rightarrow$) Let

$$\begin{aligned} M_{n+1} &\stackrel{\text{def}}{=} (M_n \uplus \sigma)|_D, \text{ where } \text{dom}(\sigma) = \mathcal{V}(M_n(g_{n+1})) \cup \mathcal{V}(M_n(\alpha_{n+1})) \\ \text{and } D &= \text{Dom}(M_n \uplus \sigma) \setminus \kappa^{-1}(q_{n+1}) \\ \text{and } M_n \uplus \sigma &\models g_{n+1} \\ (M_n \uplus \sigma)(\alpha_{n+1}) &= w_{n+1} \end{aligned}$$

We must show the existence of a substitution σ' such that

$$\begin{aligned} M'_{n+1} &\stackrel{\text{def}}{=} (M'_n \uplus \sigma')|_{D'}, \text{ where} \\ \text{dom}(\sigma') &= \mathcal{V}(M'_n(g_{n+1} \wedge \phi_{n+1})) \cup \mathcal{V}(M'_n(\alpha_{n+1})) \\ \text{and } D' &= \text{Dom}(M'_n \uplus \sigma') \setminus \kappa^{-1}(q_{n+1}) \\ \text{and } M'_n \uplus \sigma' &\models g_{n+1} \wedge \phi_{n+1} \\ (M'_n \uplus \sigma')(\alpha_{n+1}) &= w_{n+1} \end{aligned}$$

From the induction hypothesis we have $\tau_n \subseteq M'_n$ and $M_n = M'_n$. Since $\tau_{n-1} \models \phi_n$ then by applying Claim 1, we get $\tau_n \models \phi_{n+1}$. Thus we take $\sigma' = \sigma$. Hence $M'_{n+1} = M_{n+1}$ and $D' = D$. We must show that $M_n \uplus \sigma \models g_{n+1} \wedge \phi_{n+1}$. But this holds since $\tau_n \subseteq M_n$ and $\tau_n \models \phi_{n+1}$. It remains to show that $\rho_{n+1} \subseteq M'_{n+1} = M_{n+1}$. But this follows from the fact that $\tau_{n+1} \stackrel{def}{=} (\tau_n)_{|dom(\tau_n) \setminus \kappa^{-1}(q_{n+1})}$ and $M_{n+1} = (M_n \uplus \sigma)_{|D}$, where D is defined above in terms of $\kappa^{-1}(q_{n+1})$.

$\Leftarrow$) Let M'_{n+1} (as in the previous direction) and we find a substitution σ such that M_{n+1} exists. We take again $\sigma' = \sigma$, and the proof is similar to that of the previous direction.

□

□

B.2 Closure properties of GVAs

The class of GVAs with ε -transitions will be denoted by ε -GVAs. We emphasize that the ε -transitions are not guarded.

Claim 4 *For a ε -GVA $\mathcal{A}^\varepsilon$ there exists a GVA $\mathcal{A}$ (without ε -transitions) satisfying $L(\mathcal{A}) = L(\mathcal{A}^\varepsilon)$.*

Proof. The construction of a GVA out of a ε -GVA is more subtle than the construction known for FAs since we need to take into account the refreshing of the variables. We define an operator Θ that transforms a ε -GVA to an equivalent ε -GVA with strictly less ε -transitions. Thus the desired GVA without ε -transitions is the least fixed-point of Θ . Intuitively, the operator Θ eliminates all the ε -transitions which are preceded by a non ε -transition.

Assume $\mathcal{A}^\varepsilon = \langle \Sigma, \mathcal{X}, Q^\varepsilon, Q_0^\varepsilon, F^\varepsilon, \delta^\varepsilon, \kappa^\varepsilon \rangle$. Let $\mathcal{Y}(q)$ be the set of states that are reachable from state q by following an ε -transition and let $\mathcal{Y}(Q') =$

$\{\Upsilon(q) \mid q \in Q'\}$, for $Q' \subseteq Q^\varepsilon$. Let $\Theta(\mathcal{A}^\varepsilon) = \langle \Sigma, \mathcal{X}, Q, Q_0, F, \delta, \kappa \rangle$ where:

$$\begin{aligned}
Q &= Q^\varepsilon \cup (Q^\varepsilon \times Q^\varepsilon) \\
\pi_1 : \mathcal{P}(Q) &\rightarrow \mathcal{P}(Q^\varepsilon) \\
Q' &\mapsto \{p \mid (p, q) \in Q'\} \\
\pi_2 : \mathcal{P}(Q) &\rightarrow \mathcal{P}(Q^\varepsilon) \\
Q' &\mapsto \{q \mid (p, q) \in Q'\} \\
Q_0 &= Q_0^\varepsilon \cup \Upsilon(Q_0^\varepsilon) \cup \pi_1^{-1}(Q_0^\varepsilon) \\
F &= F^\varepsilon \cup \Upsilon^{-1}(F^\varepsilon) \cup \pi_2^{-1}(F^\varepsilon) \\
\delta &= \{p \xrightarrow{\alpha, g} q \in \delta^\varepsilon \mid \alpha \neq \varepsilon\} \cup \{q_1 \xrightarrow{\alpha, g} (q_2, q_3) \mid q_1 \xrightarrow{\alpha, g} q_2 \xrightarrow{\varepsilon} q_3 \in \delta^\varepsilon \mid \alpha \neq \varepsilon\} \cup \\
&\quad \{(q_1, q_2) \xrightarrow{\alpha, g} q_3 \mid q_2 \xrightarrow{\alpha, g} q_3 \in \delta^\varepsilon\} \cup \{q_1 \xrightarrow{\varepsilon} q_2 \mid \nexists q_0 \xrightarrow{\alpha, g} q_1 \text{ s.t. } \alpha \neq \varepsilon\} \\
\kappa &= \kappa^\varepsilon \cup (\pi_1^{-1} \circ \kappa^\varepsilon) \cup (\pi_2^{-1} \circ \kappa^\varepsilon)
\end{aligned}$$

In order to prove that $L(\Theta(\mathcal{A}^\varepsilon)) = L(\mathcal{A}^\varepsilon)$, it suffices to prove the following three Claims, the first one is straightforward:

Fact 1. *Every accepting run in $\mathcal{A}^\varepsilon$ that does not follow any ε -transition is still an accepting run in $\Theta(\mathcal{A}^\varepsilon)$. Conversely, every accepting run in $\Theta(\mathcal{A}^\varepsilon)$ that passes only through states in Q^ε is still an accepting run in $\mathcal{A}^\varepsilon$.*

Fact 2. *There exists a run*

$$q_0, M_0 \xrightarrow{\alpha, g} q_1, M_1 \xrightarrow{\varepsilon} q_2, M_2$$

in $\mathcal{A}^\varepsilon$ with $\alpha \neq \varepsilon$ iff there exists a run

$$q_0, M_0 \xrightarrow{\alpha, g} (q_1, q_2), M'_2$$

in $\Theta(\mathcal{A}^\varepsilon)$ such that $M_2 = M'_2$.

Proof of the Fact 2.

$\Rightarrow$) From the definition of Q and δ it follows that $(q_0, q_1) \in \delta(q_0, \alpha)$, and it remains to show that $M_2 = M'_2$. We only discuss the case when α is a letter in Σ , the case when it is a variable can be handled similarly. On the one hand, $M_2 = M_1|_{D_2}$ where $D_2 = \text{Dom}(M_1) \setminus (\kappa^\varepsilon)^{-1}(q_2)$, and $M_1 = (M_0 \uplus \gamma)|_{D_1}$ where $D_1 = \text{Dom}(M_0) \setminus \kappa^{-1}(q_1)$ and γ is a substitution such that $\gamma \models M_0(g)$. Hence $M_2 = M_0|_D$ where $D = (\text{Dom}(M_0) \uplus \text{dom}(\gamma)) \setminus ((\kappa^\varepsilon)^{-1}(q_1) \cup (\kappa^\varepsilon)^{-1}(q_2))$. On the other hand, we have $M'_2 = (M_0 \uplus \text{dom}(\gamma)) \setminus D'$, where $D' = \text{Dom}(M_0) \setminus \kappa^{-1}((q_1, q_2))$. It follows from the definition of κ , the refreshing function of $\Theta(\mathcal{A}^\varepsilon)$, that $\kappa^{-1}((q_1, q_2)) = (\kappa^\varepsilon)^{-1}(q_1) \cup (\kappa^\varepsilon)^{-1}(q_2)$. Hence, $D = D'$ and $M_2 = M'_2$.

$\Leftarrow$) This direction is proved by following the same reasoning made in the direction $(\Rightarrow)$ on the refreshing function.

This ends the proof of Fact 2. $\square$

Fact 3. *Let $q_1 \in Q^\varepsilon$ and $(q_0, q_1) \in Q$. There exists a run*

$$q_1, M_1 \xrightarrow{\alpha, g} q_2, M_2$$

in $\mathcal{A}^\varepsilon$ iff there exists a run

$$(q_0, q_1), M_1 \xrightarrow{\alpha, g} q_2, M_2$$

in $\Theta(\mathcal{A}^\varepsilon)$.

Proof of Fact 3. By checking the transition function δ . $\square$

To accomplish the proof, it remains to notice that if $q \in Q$ is such that $q \notin \pi_1^{-1}(Q^\varepsilon)$, then the outgoing transitions from q in $\mathcal{A}^\varepsilon$ are exactly the outgoing transitions from q in $\Theta(\mathcal{A}^\varepsilon)$. $\square$

Theorem 1. *GVAs are closed under concatenation, Kleene operator and intersection.*

Proof. Let $\mathcal{A}_1 = \langle \Sigma_1, \mathcal{X}_1, Q_1, q_0^1, \delta_1, F_1, \kappa_1 \rangle$ and $\mathcal{A}_2 = \langle \Sigma_2, \mathcal{X}_2, Q_2, q_0^2, \delta_2, F_2, \kappa_2 \rangle$ be two GVAs. Up to variable renaming it is sufficient to consider the closure under the above operations for two GVAs that do not share variables.

The closure under Kleene operation and concatenation is a direct consequence of the fact that GVAs with ε -transitions and GVAs recognize the same language, Claim 4 above. More precisely, the Kleene closure $\mathcal{A}_1^*$ amounts to adding an (unguarded) ε -transition between the accepting states and initial states of $\mathcal{A}_1$. And the concatenation $\mathcal{A}_1 \cdot \mathcal{A}_2$ amounts to adding an (unguarded) ε -transition between the accepting states of $\mathcal{A}_1$ and the initial states of $\mathcal{A}_2$.

The closure under intersection follows from the fact that the intersection of two GVAs $\mathcal{A}_1$ and $\mathcal{A}_2$ denoted by $\mathcal{A}_1 \cap \mathcal{A}_2$ can be defined as follows:

$$\mathcal{A}_1 \cap \mathcal{A}_2 = \langle \Sigma_1 \cup \Sigma_2, \mathcal{X}_1 \cup \mathcal{X}_2, Q_1 \times Q_2, q_0^1 \times q_0^2, \delta, F_1 \times F_2, \kappa \rangle,$$

where δ and κ are defined by:

$$\begin{cases} (q'_1, q'_2) \in \delta((q_1, q_2), (\alpha_1, (\alpha_1 = \alpha_2) \wedge g_1 \wedge g_2)) & \text{iff } q'_1 \in \delta_1(q_1, \alpha_1, g_1) \text{ and } \\ & q'_2 \in \delta_2(q_2, \alpha_2, g_2). \\ (q_1, q_2) \in \kappa(x) & \text{iff } q_1 \in \kappa_1(x) \text{ or } q_2 \in \kappa_2(x). \end{cases}$$

The proof that $L(\mathcal{A}_1) \cap L(\mathcal{A}_2) = L(\mathcal{A}_1 \cap \mathcal{A}_2)$ is straightforward. $\square$

Proposition 1. *The complement of a regular language is GVA-recognizable. That is, given a FA F there exists a GVA $\mathcal{A}$ such that $L(\mathcal{A}) = \Sigma^* \setminus L(F)$.*

Proof. The construction of $\mathcal{A}$ is similar to the one for FAs (over a finite alphabet). We assume that F is deterministic. Firstly, we make the completion of F , i.e. we construct an equivalent GVA so that for each state q of F and for each letter $l \in \Sigma$ there is a unique transition outgoing from q that reads l . Secondly, we swap the accepting and non-accepting states.

Formally, assume $F = \langle \Sigma, Q, p_0, \delta, F \rangle$, with $Q = \{q_1, \dots, q_n\}$, and define $\mathcal{A} = \langle \Sigma, \mathcal{X}, Q', p_0, \delta', F', \kappa \rangle$ by

$$\begin{cases} \mathcal{X} = \{x_1, \dots, x_n\} \\ Q' = Q \cup \{q_{i1}, q_{i2}, i = 1, \dots, n\} \\ F' = (Q \setminus F) \cup \{q_{i1}, q_{i2}, i = 1, \dots, n\} \\ \delta' = \delta \cup \left\{ q_i \xrightarrow{x_i, \wedge_j (x_i \neq a_j)} q_{i1} \mid \text{for all } a_j \text{ s.t. } q_i \xrightarrow{a_j} q_{i'} \in \delta \right\} \cup \{q_{i2} \xrightarrow{x_i} q_{i2}\} \\ \kappa(x_i) = \{q_{i2}\} \end{cases}$$

Notice that F rejects a word w iff $\mathcal{A}$ accepts w . $\square$

B.3 Decision procedures for GVAs

Theorem 2. *For GVAs, Membership is NP-complete.*

Proof. Let $\mathcal{A}$ be a GVA and $w = w_1 \dots w_n$ a word in Σ^* .

For the upper bound of the membership, a non deterministic polynomial algorithm guesses a path in $\mathcal{A}$ of length $|w|$ such that the final state is accepting; and a series of substitutions $\sigma_1, \dots, \sigma_{|w|}$, where $\sigma_i : \mathcal{X} \rightarrow \{w_j, 1 \leq j \leq |w|\}$, then checks whether the corresponding run on w is possible. The lower bound, i.e. the NP-hardness, follows from the fact that the membership problem for GVAs without guards, i.e. GVAs, is NP-complete [3, Theorem 3]. The undecidability of the universality follows from [8] since this problem is undecidable for the class of variable automata which is a subclass of GVAs. $\square$

Nonemptiness

Lemma 2. *Let $\mathcal{A} = \langle \Sigma, \mathcal{X}, Q, Q_0, F, \delta, \kappa \rangle$ be a GVA and $\mathcal{A}' = \langle \Sigma, \mathcal{X}, Q', Q'_0, F', \delta', \kappa' \rangle$ be its stretched form. Then,*

- i) *There exists a run $q_0, M_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} q_n, M_n$ in $\mathcal{A}$ if, and only if, there exists a run $(q_0, \mathcal{X}), M_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} (q_n, X_n), M_n$ in $\mathcal{A}'$ with $X_n = \mathcal{X} \setminus \text{Dom}(M_n)$.*
- ii) *A variable x is free in a state $(q, X) \in Q'$ iff $q \in X$.*

Proof.

- i) By induction on n . Since $\text{dom}(M_0) = \emptyset$ the case $n = 0$ is trivial. Assume the claim holds up to n . Let us prove the equivalence for $n + 1$.
 - $\Leftarrow$) Since $(q_{n+1}, X_{n+1}) \in \delta'((q_n, X_n), \alpha_{n+1}, g_{n+1})$ by induction $X_n = \mathcal{X} \setminus \text{Dom}(M_n)$. Thus $q_{n+1} \in \delta(q_n, \alpha_{n+1}, g_{n+1})$. The substitution M_{n+1} obtained is as expected.
 - $\Rightarrow$) Assume a transition $q_n, M_n \xrightarrow{\alpha_{n+1}, g_{n+1}} q_{n+1}, M_{n+1}$. From the definition of δ' it follows that $(q_{n+1}, X_{n+1}) \in \delta'((q_n, X_n), \alpha_{n+1}, g_{n+1})$.
- ii) Immediate from Item i) since for every configuration $((q, X), M)$ we have that $x \in X$ iff $x \notin \text{Dom}(M)$.

□

Lemma 3. *There exists a run $q_0, M_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} q_n, M_n$ in $\mathcal{A}$ if, and only if, there exists a run $(q_0, \text{true}), M_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} (q_n, \phi_n), M_n$ in $\mathcal{A}_{/\approx}$ such that $M_i \models \phi_i$, for all $i = 0, \dots, n$.*

Proof. By induction on n . Since $\text{dom}(M_0) = \emptyset$ and $g_0 = \text{true}$ the case $n = 0$ is trivial. Assume the claim holds up to n . Let us prove the equivalence for $n + 1$.

- $\Leftarrow$) Since $(q_{n+1}, \phi_{n+1}) \in \delta'((q_n, \phi_n), \alpha_{n+1}, g_{n+1})$ then $q_{n+1} \in \delta(q_n, \alpha_{n+1}, g_{n+1})$ where $\phi_{n+1} = (g_{n+1} \wedge \phi_n)^{\kappa^{-1}(q_{n+1})}$. Thus $q_{n+1} \in \delta(q_n, \alpha_{n+1}, g_{n+1})$. From the induction hypothesis we have

$$M_n \models \phi_n, \tag{2}$$

and we know that there exists a substitution σ such that $\alpha_{n+1} \in \text{Dom}(\sigma)$ and

$$\begin{cases} M_{n+1} = (\sigma \uplus M_n)|_D, \text{ and} \\ \sigma \uplus M_n \models g_{n+1} \end{cases} \tag{3}$$

where $D = \text{Dom}(\sigma \uplus M_n) \setminus \kappa^{-1}(q_{n+1})$. We must show that

$$M_{n+1} \models (g_{n+1} \wedge \phi_n)^{\kappa^{-1}(q_{n+1})}$$

Since $M_{n+1} = (\sigma \uplus (M_n))|_D$ (Eq. 3), we must show that

$$(\sigma \uplus (M_n))|_D \models g_{n+1}^{\kappa^{-1}(q_{n+1})} \wedge \phi_n^{\kappa^{-1}(q_{n+1})} \quad (4)$$

On the one hand, it follows from 3 that $(\sigma \uplus (M_n))|_D \models g_{n+1}^{\kappa^{-1}(q_{n+1})}$. On the other hand, from 2 we get $(\sigma \uplus (M_n))|_D \models \phi_n^{\kappa^{-1}(q_{n+1})}$. And we are done.

$\Rightarrow$) Let $q_{n+1} \in \delta(q_n, \alpha_{n+1}, g_{n+1})$. Hence $(q_{n+1}, \phi_{n+1}) \in \delta'((q_n, \phi_n), \alpha_{n+1}, g_{n+1})$ then $q_{n+1} \in \delta(q_n, \alpha_{n+1}, g_{n+1})$ where $\phi_{n+1} = (g_{n+1} \wedge \phi_n)^{\kappa^{-1}(q_{n+1})}$. By using similar argument, the claim $M_{n+1} \models \phi_{n+1}$ holds. $\square$

Fact 3 If $q_0, M_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} q_n, M_n$ is a run in $\mathcal{A}$, where $n \geq 1$, then $M_i \vdash g_{i+1}$ for all $i = 0, \dots, n-1$.

Proof. This is an immediate consequence of the definition of the run, since there are substitutions γ_i such that $\gamma_i \uplus M_i \models g_{i+1}$. $\square$

Lemma 4. Let $\mathcal{A}$ be a GVA in the normal form. Let

$$\pi = (q_0, X_0, \phi_0) \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} (q_n, X_n, \phi_n)$$

be a path in $\mathcal{A}$ with $\phi_0 = \text{true}$ and $q_0 \in Q_0$. Then, there is a run over π iff $\vdash \phi_n$ and $\vdash \phi_i \wedge g_{i+1}$, for all $i = 0, \dots, n-1$.

Proof. From Lemma 3 and Fact 3 it follows that there is a run

$$\pi^r = (q_0, X_0, \phi_0), M_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} (q_n, X_n, \phi_n), M_n$$

iff $M_i \models \phi_i$ for all $i = 0, \dots, n$, and $M_j \vdash g_{j+1}$ for all $j = 0, \dots, n-1$. Therefore $M_i \vdash \phi_i$, hence $M_i \vdash \phi_i \wedge g_{i+1}$. Thus we get $\vdash \phi_i \wedge g_{i+1}$ for all $i = 0, \dots, n-1$, and $\vdash \phi_n$. $\square$

Corollary 2. Let $\mathcal{A}$ be a GVA in the normal form. Let $\pi_1 \circ \pi_2 \circ \pi_3$ be a path in $\mathcal{A}$ with $\eta_f(\pi_1) = \eta_0(\pi_2) = \eta_f(\pi_2) = \eta_0(\pi_3)$. If there is an accepting run in $\mathcal{A}$ over $\pi_1 \circ \pi_2 \circ \pi_3$ then there is an accepting run over $\pi_1 \circ \pi_3$ as well.

Proof. Let

$$\begin{aligned}\pi_1 &= (q_0, X_0, \phi_0) \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} (q_n, X_n, \phi_n) \\ \pi_2 &= (q_n, X_n, \phi_n) \xrightarrow{\alpha_{n+1}, g_{n+1}} \dots \xrightarrow{\alpha_m, g_m} (q_m, X_m, \phi_m) = (q_n, X_n, \phi_n) \\ \pi_3 &= (q_m, X_m, \phi_m) \xrightarrow{\alpha_{m+1}, g_{m+1}} \dots \xrightarrow{\alpha_r, g_r} (q_r, X_r, \phi_r)\end{aligned}$$

be paths in $\mathcal{A}$ and assume the existence of an accepting run over $\pi_1 \circ \pi_2 \circ \pi_3$ then it follows from Lemma 4 that $\vdash \phi_r$ and $\vdash \phi_i \wedge g_{i+1}$, for $i = 0, \dots, r-1$. Hence $\vdash \phi_i \wedge g_{i+1}$, for $i \in [0, \dots, n-1] \cup [m, r-1]$. That is, there is an accepting run over the path $\pi_1 \circ \pi_3$. $\square$

C Proofs for Section 5

The definition of simulation preorder for GVAs follows.

Definition 11. Let $\mathcal{A}_1 = \langle \Sigma, \mathcal{X}_1, Q_1, q_0^1, \tau_1, \delta_1, F_1, \kappa_1 \rangle$ and $\mathcal{A}_2 = \langle \Sigma, \mathcal{X}_2, Q_2, q_0^2, \tau_2, \delta_2, F_2, \kappa_2 \rangle$ be two GVAs where $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$. A simulation of $\mathcal{A}_1$ by $\mathcal{A}_2$ is a relation $\leq \subseteq (\zeta_{\mathcal{X}_1, \Sigma} \times Q_1) \times (\zeta_{\mathcal{X}_2, \Sigma} \times Q_2)$ such that:

- $(\tau_1, q_0^1) \leq (\tau_2, q_0^2)$.
- if $(\sigma_1, q_1) \leq (\sigma_2, q_2)$ and if $(\sigma_1, q_1) \xrightarrow{a} (\sigma'_1, q'_1)$ for $a \in \Sigma$ then there exists a state $q'_2 \in Q_2$ such that $(\sigma_2, q_2) \xrightarrow{a} (\sigma'_2, q'_2)$ and $(\sigma'_1, q'_1) \leq (\sigma'_2, q'_2)$.

The claims in the following Lemma are not hard to prove. They will be used in the proofs of the main claims.

Lemma 4. Let $C \subseteq \Sigma$ be a finite set of letters, $\bar{\sigma}$ and σ two substitutions, x , and a a letter in C . The following hold.

1. If $\bar{\sigma} \bowtie_C \sigma$ then $|\text{codom}(\bar{\sigma})| = |\text{codom}(\sigma)|$.
2. If $\bar{\sigma} \bowtie_C \sigma$ and $D \subseteq \text{Dom}(\sigma)$ then $\bar{\sigma}|_D \bowtie_C \sigma|_D$,
3. If $(\bar{\sigma}_1 \uplus \bar{\sigma}_2) \bowtie (\sigma_1 \uplus \sigma_2)$ with $\text{dom}(\bar{\sigma}_i) = \text{dom}(\sigma_i)$, then $\bar{\sigma}_i \bowtie \sigma_i$, for $i = 1, 2$.
4. If $\bar{\sigma} \bowtie_C \sigma$ and γ is a substitution with $\text{dom}(\gamma) \cap \text{dom}(\sigma) = \emptyset$, then $\bar{\sigma} \uplus \gamma \bowtie_C \sigma \uplus \gamma$.
5. If $\bar{\sigma} \bowtie_C \sigma$ with $\bar{\sigma}(y) = \bar{a}$ and $\sigma(y) = a$ for some variable y , and $x \notin \text{dom}(\sigma)$ then $\bar{\sigma} \uplus \{x \mapsto \bar{a}\} \bowtie_C \sigma \uplus \{x \mapsto a\}$.
6. If $\bar{\sigma} \bowtie_C \sigma$ and $\bar{a} \notin C \cup \text{codom}(\bar{\sigma})$ and $a \notin C \cup \text{codom}(\sigma)$ and $x \notin \text{dom}(\sigma)$ then $\bar{\sigma} \uplus \{x \mapsto \bar{a}\} \bowtie_C \sigma \uplus \{x \mapsto a\}$.

Notice that the opposite direction of the Item 3 of Lemma 4 does not hold in general.

Lemma 5. *Let σ and $\bar{\sigma}$ be two substitution, where $\sigma \bowtie_C \bar{\sigma}$, and let g be a guard such that $\Sigma_g \subseteq C$. Then, $\sigma \models g$ iff $\bar{\sigma} \models g$.*

Proof. By induction on the structure of g in both directions.

$\Rightarrow$) If $g = (\alpha = x)$, where $\alpha \in \Sigma$ and $x \in \mathcal{X}$, then $\sigma(x) \in C$, hence $\bar{\sigma}(x) \in C$. Therefore $\sigma(x) = \bar{\sigma}(x) = \alpha$. If $g = (x = y)$, where $x, y \in \mathcal{X}$, then from the definition of $\bowtie_C$ we have that $\sigma(x) = \sigma(y)$ iff $\bar{\sigma}(x) = \bar{\sigma}(y)$. Thus the claim holds. The case when $g = g_1 \wedge g_2$ follows from a direct application of the induction hypothesis.

$\Leftarrow$) This direction follows from the fact that $\sigma \bowtie_C \bar{\sigma}$ iff $\bar{\sigma} \bowtie_C \sigma$. □

Corollary 3. *Let σ and $\bar{\sigma}$ be two substitution, where $\sigma \bowtie_C \bar{\sigma}$, and let g be a guard such that $\Sigma_g \subseteq C$. Then, $\sigma \vdash g$ iff $\bar{\sigma} \vdash g$.*

Proof.

$\Rightarrow$) We show that if there exists a substitution γ such that $\text{dom}(\gamma) = \mathcal{V}(g) \setminus \text{dom}(\sigma)$ and $\sigma \models \gamma(g)$, then $\bar{\sigma} \models \gamma(g)$. But this follows from Lemma 5. 5.

$\Leftarrow$) This direction follows from the fact that $\bowtie_C$ is symmetric relation. □

Corollary 4. *Let $\sigma, \bar{\sigma}, \gamma, \bar{\gamma}$ be substitutions, where $\text{dom}(\gamma) \cap \text{dom}(\sigma) = \emptyset$ and $\text{dom}(\bar{\gamma}) \cap \text{dom}(\bar{\sigma}) = \emptyset$. Let g be a guard such that $\Sigma_g \subseteq C$. If $\sigma \uplus \gamma \bowtie_C \bar{\sigma} \uplus \bar{\gamma}$ then we have that $\gamma \models \sigma(g)$ iff $\bar{\gamma} \models \bar{\sigma}(g)$.*

In what follows we let S_1, S_2 be two (possibly infinite) sets of letters with $|S_1 \setminus S_2| > |\mathcal{X}|$ and $|S_2 \setminus S_1| > |\mathcal{X}|$ and $S_1 \cap S_2 \neq \emptyset$. Let $C = S_1 \cap S_2$.

In order to relate the instantiations of the variables in the game $\mathcal{G}$ to the instantiation of the variables in the game $\overline{\mathcal{G}}$, we need to introduce the function $\Theta_C^{S_1, S_2} : \xi_{\mathcal{X}, S_1} \times \xi_{\mathcal{X}, S_1} \times \xi_{\mathcal{X}, S_2} \rightarrow \xi_{\mathcal{X}, S_2}$, that, given three substitutions M, γ and M' such that $\text{dom}(M) \cap \text{dom}(\gamma) = \emptyset$ and $M \bowtie_C M'$, constructs a substitution $\gamma' = \Theta_C^{S_1, S_2}(M, \gamma, M')$ such that $M \uplus \gamma \bowtie_C M' \uplus \gamma'$.

Definition 12. *We define the functions*

$$\Theta_C^{S_1, S_2}, \bar{\Theta}_C^{S_1, S_2} : \xi_{\mathcal{X}, S_1} \times \xi_{\mathcal{X}, S_1} \times \xi_{\mathcal{X}, S_2} \rightarrow \xi_{\mathcal{X}, S_2}$$

as follows. Let $M_1, \gamma_1 \in \xi_{\mathcal{X}, S_1}$, $M_2 \in \xi_{\mathcal{X}, S_2}$. Then, $\Theta_C^{S_1, S_2}(M_1, \gamma_1, M_2)$ is defined only when $|\text{dom}(\gamma_1)| = 1$ and $\text{dom}(\gamma_1) \cap \text{dom}(M_1) = \emptyset$ and

$M_1 \bowtie_C M_2$ by:

$$\Theta_C^{S_1, S_2}(M_1, \gamma_1, M_2) = \begin{cases} \gamma_1 & \text{if } \gamma_1(x) \in C \\ \{x \mapsto M_2(y)\} & \text{if } \gamma_1(x) \in \text{codom}(M_1) \setminus C \text{ and} \\ & M_1(y) = \gamma_1(x), y \in \mathcal{X} \\ \{x \mapsto \text{get}(S_2 \setminus \text{codom}(M_2))\} & \text{if } \gamma_1(x) \in S_1 \setminus (C \cup \text{codom}(M_1)) \end{cases}$$

where $\text{dom}(\gamma_1) = \{x\}$.

And $\Theta_C^{S_1, S_2}(M_1, \gamma_1, M_2)$ is defined only when $\text{dom}(\gamma_1) \cap \text{dom}(M_1) = \emptyset$ by:

$$\Theta_C^{S_1, S_2}(M_1, \gamma_1, M_2) = \begin{cases} \Theta_C^{S_1, S_2}(M_1, \gamma_1, M_2) & \text{if } |\gamma_1| = 1 \\ \gamma'_2 \uplus \Theta_C^{S_1, S_2}(M_1 \uplus \gamma'_1, \gamma''_1, M_2 \uplus \gamma'_2) & \text{if } |\gamma_1| \geq 2, \gamma_1 = \gamma'_1 \uplus \gamma''_1 \text{ and } |\gamma'_1| = 1, \\ & \text{where } \gamma'_2 = \Theta_C^{S_1, S_2}(M_1, \gamma'_1, M_2) \end{cases}$$

Lemma 6. Let $M_1, \gamma_1 \in \xi_{\mathcal{X}, S_1}$ and $M_2 \in \xi_{\mathcal{X}, S_2}$ be substitutions with $\text{dom}(M_1) \cap \text{dom}(\gamma_1) = \emptyset$ and $M_1 \bowtie_C M_2$. We have that

$$(M_1 \uplus \gamma_1) \bowtie_C (M_2 \uplus \Theta_C^{S_1, S_2}(M_1, \gamma_1, M_2))$$

Proof. By induction on $|\text{dom}(\gamma_1)| = 1$.

Base Case. If $|\text{dom}(\gamma_1)| = 1$ then assume $\text{dom}(\gamma_1) = \{x\}$ and let $\gamma_2 = \Theta_C^{S_1, S_2}(M_1, \gamma_1, M_2)$. We distinguish three cases depending on $\gamma_1(x)$.

- If $\gamma_1(x) \in C$ then it follows from the definition of $\Theta_C^{S_1, S_2}$ that $\gamma_2 = \gamma_1$. From the Item 4 of Lemma 4 we get $M_1 \uplus \gamma_1 \bowtie_C M_2 \uplus \gamma_1$.
- If $\gamma_1(x) \in \text{codom}(M_1) \setminus C$ then in this case we recall that $\gamma_2 = \{x \mapsto M_2(y)\}$ where $M_1(y) = \gamma_1(x)$ for some variable $y \in \mathcal{X}$. The claim that $(M_1 \uplus \{x \mapsto \gamma_1(x)\}) \bowtie_C M_2 \uplus \{x \mapsto M_2(y)\}$ follows from the Item 5 of Lemma 4.
- Otherwise, i.e. if $\gamma_1(x) \in S_1 \setminus (C \cup \text{codom}(M_1))$ then the claim that $(M_1 \uplus \{x \mapsto \gamma_1(x)\}) \bowtie_C (M_2 \uplus \{x \mapsto \text{get}(S_2 \setminus \text{codom}(M_2))\})$ follows from the Item 6 of Lemma 4.

Induction Case. Assume $\gamma_1 = \gamma'_1 \uplus \gamma''_1$ with $|\gamma'_1| = 1$. Let

$$\begin{cases} \gamma'_2 = \Theta_C^{S_1, S_2}(M_1, \gamma'_1, M_2), \text{ and} \\ \gamma''_2 = \Theta_C^{S_1, S_2}(M_1 \uplus \gamma'_1, \gamma''_1, M_2 \uplus \gamma'_2) \text{ and} \\ \gamma_2 = \Theta_C^{S_1, S_2}(M_1, \gamma_1, M_2) = \gamma'_2 \uplus \gamma''_2 \end{cases}$$

From the induction hypothesis it follows that

$$\begin{cases} M_1 \uplus \gamma'_1 & \bowtie_C M_2 \uplus \gamma'_2 \\ (M_1 \uplus \gamma'_1) \uplus \gamma''_1 & \bowtie_C (M_2 \uplus \gamma'_2) \uplus \gamma''_2 \end{cases}$$

Therefore

$$M_1 \uplus \gamma_1 \bowtie_C M_2 \uplus \gamma_2$$

□

Lemma 7. *Let $\sigma_1, \sigma_2, \sigma_3, \gamma_2 \in \xi_{\mathcal{X}, S_1}$ and $\alpha, \beta \in C \cup \mathcal{X}$ and $\sigma'_1, \sigma'_2, \sigma'_3 \in \xi_{\mathcal{X}, S_2}$ and $g_2 \in G$ be such that*

$$\begin{cases} \gamma_2(\sigma_2(\beta)) = \sigma_3(\alpha) \\ \sigma_3 \uplus \sigma_2 \bowtie_C \sigma'_3 \uplus \sigma'_2 \\ \sigma_1 \bowtie_C \sigma'_1 \\ \gamma_2 \models \sigma_2(g_2) \\ \text{dom}(\sigma_1) \subseteq \text{dom}(\sigma_3) \\ \text{dom}(\sigma'_1) \subseteq \text{dom}(\sigma'_3) \end{cases} \quad (5)$$

There exists a function $\Xi_C^{S_1, S_2}$

$$\begin{aligned} \Xi_C^{S_1, S_2} : \quad & \xi_{\mathcal{X}, S_1}^5 \times (C \cup \mathcal{X})^2 \times G_C \times \xi_{\mathcal{X}, S_2}^3 \rightarrow \xi_{\mathcal{X}, S_2} \\ & (\sigma_1, \sigma_2, \sigma_3, \gamma_2, \alpha, \beta, g_2, \sigma'_1, \sigma'_2, \sigma'_3) \mapsto \gamma'_2 \end{aligned}$$

which is defined only if Eq (5) holds and satisfies the following:

$$\gamma'_2(\sigma'_2(\beta)) = \sigma'_3(\alpha) \quad (\text{A.1})$$

$$(\sigma_1 \uplus \gamma_1) \uplus (\sigma_2 \uplus \gamma_2) \bowtie_C (\sigma'_1 \uplus \gamma'_1) \uplus (\sigma'_2 \uplus \gamma'_2) \quad (\text{A.2})$$

$$\gamma'_2 \models \sigma'_2(g_2) \quad (\text{A.3})$$

Proof. The construction of $\Xi_C^{S_1, S_2}$ depends on $\sigma_3(\alpha)$

- I.) If $\sigma_3(\alpha) \in C$, then in this case we have $\sigma_3(\alpha) = \sigma'_3(\alpha) \in C$. Hence $\sigma'_2 = \sigma_2$. Thus (A.1) holds. Furthermore we let:

$$\gamma'_2 = \Theta_C^{S_1, S_2}(\sigma_3 \uplus \sigma_2, \gamma_2, \sigma'_3 \uplus \sigma'_2)$$

From Lemma 6 it follows that

$$\sigma_3 \uplus \sigma_2 \uplus \gamma_1 \uplus \gamma_2 \bowtie_C \sigma'_3 \uplus \sigma'_2 \uplus \gamma'_1 \uplus \gamma'_2$$

From Eq (5) we have $\text{dom}(\sigma_1) \subseteq \text{dom}(\sigma_3)$ and $\text{dom}(\sigma'_1) \subseteq \text{dom}(\sigma'_3)$ and $\text{dom}(\sigma_1) = \text{dom}(\sigma'_1)$, hence it follows from Item 2 of Lemma 4 that

$$\sigma_1 \uplus \sigma_2 \uplus \gamma_2 \bowtie_C \sigma'_1 \uplus \sigma'_2 \uplus \gamma'_2$$

Therefore (A.2) holds. Finally (A.3) follows from Corollary 4.

- II.) If $\sigma_3(\alpha) \in S_1 \setminus C$, then α must be a variable, say $y_1 \in \mathcal{X}$. We distinguish two cases depending on $\sigma_2(\beta)$.
- i.) If $\sigma_2(\beta)$ is a letter then in this case $\sigma_2(\beta) = \sigma_3(\alpha)$, and we let

$$\gamma'_2 = \Theta_C^{S_1, S_2}(\sigma_3 \uplus \sigma_2 \uplus \gamma_1, \gamma_2, \sigma'_3 \uplus \sigma'_2 \uplus \gamma'_1)$$

And we must show that $\sigma'_3(\alpha) = \sigma'_2(\beta)$. Notice that β must be a variable, say $y_2 \in \mathcal{X}$. On the one hand we have that $\{y_1 \mapsto \sigma_3(\alpha), y_2 \mapsto \sigma_2(\beta)\} \subseteq \sigma_3 \uplus \sigma_2$ and $\{y_1 \mapsto \sigma'_3(\alpha), y_2 \mapsto \sigma'_2(\beta)\} \subseteq \sigma'_3 \uplus \sigma'_2$. On the other hand, we have that $\sigma_3(\alpha) = \sigma_2(\beta)$ and $\sigma_3 \uplus \sigma_2 \bowtie_C \sigma'_3 \uplus \sigma'_2$. Therefore $\sigma'_3(\alpha) = \sigma'_2(\beta)$, thus (A.1), (A.2) and (A.3) hold.

- ii.) If $\sigma_2(\beta)$ is a variable, say $y_2 \in \mathcal{X}$, then $\sigma'_2(\beta) = \sigma_2(\beta) = \beta = y_2$, since $\sigma'_2 \bowtie \sigma_2$. In this case we have $\{y_2 \mapsto \sigma_3(\alpha)\} \subseteq \gamma_2$. Thus we let

$$\gamma'_2 = \Theta_C^{S_1, S_2}(\sigma_3 \uplus \sigma_2 \uplus \{y_1 \mapsto \sigma_3(\alpha)\}, \gamma_2, \sigma'_3 \uplus \sigma'_2 \uplus \{y_1 \mapsto \sigma'_3(\alpha)\})$$

And Eqs (A.1), ..., (A.3) hold.

- III.) If $\sigma_3(\alpha)$ is a variable, say $x_1 \in \mathcal{X}$, then $\sigma'_3(\alpha) = \sigma_3(\alpha) = \alpha = x_1$. We distinguish two cases depending on the nature of $\sigma_2(\beta)$.

- I.) If $\sigma_2(\beta)$ is a letter then $\sigma'_2(\beta)$ is a letter as well since $\sigma'_2 \bowtie \sigma_2$. This case is dual w.r.t. case II.ii).

- II.) If $\sigma_2(\beta)$ is a variable, say $y_2 \in \mathcal{X}$, then $\sigma_2(\beta) = \sigma'_2(\beta) = \beta = y_2$ since $\sigma'_2 \bowtie \sigma_2$. In this case we let $\gamma'_2 = \Theta_C^{S_1, S_2}(\sigma_1 \uplus \sigma_2, \gamma_2, \sigma'_1 \uplus \sigma'_2)$.

□

By using the functions Θ and ξ and their respective properties stated in Lemmas 6 and 7 we are ready to prove that the games $\mathcal{G}$ and $\overline{\mathcal{G}}$ are equivalent:

Proposition 2. *Let $\mathcal{A}_1 = \langle \Sigma, \mathcal{X}_1, Q_1, q_0^1, \tau_1, \delta_1, F_1, \kappa_1 \rangle$ and $\mathcal{A}_2 = \langle \Sigma, \mathcal{X}_2, Q_2, q_0^2, \tau_2, \delta_2, F_2, \kappa_2 \rangle$ be two GVAs. Then, *Eloise* has a winning strategy in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ iff she has a winning strategy in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$.*

Proof. Up to variables renaming, we can assume that $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$. For the direction " $\Rightarrow$ " we show that out of a winning strategy of **Eloise** in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ we construct a winning strategy for her in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$. For this purpose, we shall show that each move of **Abelard** in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ can be mapped to an **Abelard** move in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$, and **Eloise** response in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ can be actually mapped to an **Eloise** move in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$. This mapping defines a relation $\mathcal{R}$ ³ between the positions of $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ and the positions of $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ as follows:

$$\mathcal{R} \subseteq \text{Pos}_E(\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)) \times \text{Pos}_E(\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)) \cup \\ \text{Pos}_A(\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)) \times \text{Pos}_A(\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2))$$

Furthermore, we impose that the following invariant holds:

$$\text{If } (\bar{\wp}, \wp) \in \mathcal{R} \text{ then } \bar{\wp} \bowtie_C \wp, \quad (\text{Inv-}\bowtie)$$

where $C = \Sigma_{\mathcal{A}_1} \cup \Sigma_{\mathcal{A}_2} \cup \text{codom}(\tau_1) \cup \text{codom}(\tau_2)$. We recall that the variables in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ are instantiated from the set of letters $C_0 = C \cup (\mathcal{X}_1 \times \mathcal{X}_2) \cup (\mathcal{X}_2 \times \mathcal{X}_1)$. The proof is by induction on n , the number of the moves made in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ plus the number of moves made in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$. The base case, i.e. when $n = 0$, trivially holds since the starting position of $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ and of $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ is $((\tau_1, q_0^1), (\tau_2, q_0^2))_{\mathbf{A}}$.

For the induction case let $(\bar{\wp}_n, \wp_n) \in \mathcal{R}$. We consider two possibilities: when $\bar{\wp}_n$ and $\wp_n$ are both **Abelard** positions and when they are both **Eloise** positions.

- i) Consider the first possibility and an **Abelard** move $\bar{m} = \bar{\wp}_n \rightarrow \bar{\wp}_{n+1}$ in $\overline{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$. In this case we have $\bar{m} \in M_A$ and $\bar{m}$ is of the form:

$$\bar{m} = ((\bar{\sigma}_1, q_1), (\bar{\sigma}_2, q_2))_{\mathbf{A}} \rightarrow (((\bar{\sigma}_1 \uplus \bar{\gamma})|_D, q'_1), (\bar{\sigma}_2, q_2), (\bar{\sigma}_1 \uplus \bar{\gamma}, g_1))_{\mathbf{E}} \\ | \quad q'_1 \in \delta_1(q_1, \alpha, g_1) \\ \text{and } D = \text{Dom}(\bar{\sigma}_1 \uplus \bar{\gamma}) \setminus \kappa_1^{-1}(q'_1) \\ \text{and } \bar{\sigma}_1 \uplus \bar{\gamma} \vdash g_1 \\ \text{and } \bar{\gamma} : \mathcal{V}(\bar{\sigma}_1(g_1)) \setminus \mathcal{V}(\bar{\sigma}_1(\alpha)) \rightarrow C_0$$

From the induction hypothesis we have $\bar{\wp}_n \bowtie_C \wp_n$. Hence $\wp_n = ((\sigma_1, q_1), (\sigma_2, q_2))_{\mathbf{A}}$, for two substitutions σ_1, σ_2 where $(\bar{\sigma}_1 \uplus \bar{\sigma}_2) \bowtie_C (\sigma_1 \uplus \sigma_2)$. Thus **Abelard** move in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$ is

$$\bar{m} = ((\sigma_1, q_1), (\sigma_2, q_2))_{\mathbf{A}} \rightarrow (((\sigma_1 \uplus \gamma)|_D, q'_1), (\sigma_2, q_2), (\sigma_1 \uplus \gamma))_{\mathbf{E}}$$

³ More precisely, if $(\bar{\wp}, \wp) \in \mathcal{R}$, and the move $\bar{\wp} \xrightarrow{\bar{\mathcal{G}}} \bar{\wp}'$ is mapped to $\wp \xrightarrow{\mathcal{G}} \wp'$, or $\wp \xrightarrow{\mathcal{G}} \wp'$ is mapped to $\bar{\wp} \xrightarrow{\bar{\mathcal{G}}} \bar{\wp}'$, then $(\bar{\wp}', \wp') \in \mathcal{R}$

where $\gamma : \mathcal{V}(\sigma_1(g_1)) \setminus \mathcal{V}(\alpha) \rightarrow \Sigma$ is defined by

$$\gamma = \Theta_C^{C_0, \Sigma}(\bar{\sigma}_1 \uplus \bar{\sigma}_2, \bar{\gamma}, \sigma_1 \uplus \sigma_2). \quad (6)$$

Indeed, notice that since $\bar{\sigma}_1 \bowtie_C \sigma_1$ then $\text{dom}(\bar{\gamma}) = \text{dom}(\gamma)$. Furthermore, we must show that

$$\sigma_1 \uplus \gamma \vdash g_1 \quad (7)$$

and that the invariant [\(Inv- \$\bowtie\$ \)](#) is maintained, i.e. to show that $\bar{\wp}_{n+1} \bowtie_C \wp_{n+1}$, that is to show that:

$$((\bar{\sigma}_1 \uplus \bar{\gamma})|_D \cup (\bar{\sigma}_1 \uplus \bar{\gamma})) \uplus \bar{\sigma}_2 \bowtie_C ((\sigma_1 \uplus \gamma)|_D \cup (\sigma_1 \uplus \gamma)) \uplus \sigma_2. \quad (8)$$

From the definition of γ in Eq (6) and by applying Lemma 6 we get: $(\bar{\sigma}_1 \uplus \bar{\sigma}_2) \uplus \bar{\gamma} \bowtie_C (\sigma_1 \uplus \sigma_2) \uplus \gamma$. Therefore,

$$(\bar{\sigma}_1 \uplus \bar{\gamma}) \uplus \bar{\sigma}_2 \bowtie_C (\sigma_1 \uplus \gamma) \uplus \sigma_2 \quad (9)$$

On the one hand, it follows from the Item 3 of Lemma 4 that $(\bar{\sigma}_1 \uplus \bar{\gamma}) \bowtie_C (\sigma_1 \uplus \gamma)$. Since we already have $\bar{\sigma}_1 \uplus \bar{\gamma} \vdash g_1$, then it follows from Corollary 3 that $\sigma_1 \uplus \gamma \vdash g_1$. Thus Eq (7) is proved. On the other hand, since $M|_D \subseteq M$ for any substitution M and any $D \subseteq \text{dom}(M)$, then Eq (8) follows from Eq (9).

- ii) Secondly, we consider the possibility when both $\bar{\wp}_n$ and $\wp_n$ are **Eloise** positions. We consider an **Eloise** move $m = \wp_n \rightarrow \wp_{n+1}$ in $\mathcal{G}(\mathcal{A}_1, \mathcal{A}_2)$, and we describe the corresponding **Eloise** move in $\bar{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$. In this case we have $m \in M_E$, and m is of the form:

$$\begin{aligned} m &= ((\sigma_1, q_1), (\sigma_2, q_2), (\sigma_3, \alpha))_E \rightarrow ((\sigma_1, q_1), ((\sigma_2 \uplus \gamma_2)|_{D_2}, q'_2))_A \\ &\quad | \quad q'_2 \in \delta_2(q_2, \beta, g_2) \\ &\quad \text{and } D_2 = \text{Dom}(\sigma_2 \uplus \gamma_2) \setminus \kappa_2^{-1}(q'_2) \\ &\quad \text{and } \gamma_1(\sigma_3(\alpha)) = \gamma_2(\sigma_2(\beta)) \\ &\quad \text{and } \gamma_2 \models \sigma_2(g_2) \\ &\quad \text{and } \gamma_2 : \mathcal{V}(\sigma_2(\beta)) \cup \mathcal{V}(\sigma_2(g_2)) \rightarrow \Sigma \end{aligned}$$

From the induction hypothesis we have that $\bar{\wp}_n \bowtie_C \wp_n$, therefore $\bar{\wp}_n = ((\bar{\sigma}_1, q_1), (\bar{\sigma}_2, q_2), (\bar{\sigma}_3, \alpha, g_1))_E$, for substitutions $\bar{\sigma}_1, \bar{\sigma}_2, \bar{\sigma}_3$, such that $((\bar{\sigma}_1 \cup \bar{\sigma}_3) \uplus \bar{\sigma}_2) \bowtie_C ((\sigma_1 \cup \sigma_3) \uplus \sigma_2)$.

The corresponding move $\bar{m}$ in $\bar{\mathcal{G}}(\mathcal{A}_1, \mathcal{A}_2)$ is:

$$\begin{aligned}\bar{m} &= ((\bar{\sigma}_1, q_1), (\bar{\sigma}_2, q_2), (\bar{\sigma}_3, \alpha))_{\mathbf{E}} \rightarrow ((\bar{\sigma}_1, q_1), ((\bar{\sigma}_2 \uplus \bar{\gamma}_2)_{|D_2}, q'_2))_{\mathbf{A}} \\ &\quad \text{and } \bar{\gamma}_1(\bar{\sigma}_3(\alpha)) = \bar{\gamma}_2(\bar{\sigma}_2(\beta)) \\ &\quad \text{and } \bar{\gamma}_2 \models \bar{\sigma}_2(g_2) \\ &\quad \text{and } \bar{\gamma}_2 : \mathcal{V}(\bar{\sigma}_2(\beta)) \cup \mathcal{V}(\bar{\sigma}_2(g_2)) \rightarrow C_0\end{aligned}$$

where $\bar{\gamma}_2$ is defined by

$$\bar{\gamma}_2 = \Xi_C^{\Sigma, C_0}(\sigma_1, \sigma_2, \sigma_3, \gamma_2, \alpha, \beta, g_2, \bar{\sigma}_1, \bar{\sigma}_2, \bar{\sigma}_3)$$

From Eq (A.2) of Lemma 7 we get

$$\sigma_1 \uplus \sigma_2 \uplus \gamma_2 \bowtie_C \bar{\sigma}_1 \uplus \bar{\sigma}_2 \uplus \bar{\gamma}_2$$

From the Item 2 of Lemma 4 it follows that the invariant is maintained, i.e.

$$\sigma_1 \uplus (\sigma_2 \uplus \gamma_2)_{|D_2} \bowtie_C \bar{\sigma}_1 \uplus (\bar{\sigma}_2 \uplus \bar{\gamma}_2)_{|D_2}$$

The proof of the direction " $\Leftarrow$ " is dual w.r.t. the proof of the direction " $\Rightarrow$ ". That is, it can be obtained by replacing **Eloise** by **Abelard**, and **Abelard** by **Eloise** and keeping the same instantiation strategy and the definition of the $\bowtie$ -coherence. This ends the proof of the Proposition. $\square$

D Proofs and definitions for Section 6

D.1 GVAs vs. variable automata of [8]

In this subsection we show that for every variable automata we can construct a GVA recognizing the same language. Firstly, we recall the definition of variable automata of Grumberg et al. [8].

Definition 13. *A variable automaton is a tuple $\mathcal{B} = \langle \Sigma, \mathcal{X} \cup \{y\}, B \rangle$, where Σ is an infinite set of letters, $\mathcal{X} \cup \{y\}$ is a finite set of variables, and $B = \langle \Sigma_B \cup \mathcal{X} \cup \{y\}, Q, Q_0, \delta, F \rangle$ is a finite automaton, called the pattern automaton, with the finite set $\Sigma_B \subset \Sigma$.*

The main idea is that the variables in $\mathcal{X} \cup \{y\}$ range over $\Sigma \setminus \Sigma_B$, and the instantiations of y differ from the instantiations of the variables in $\mathcal{X}$. The formal definition of languages recognized by variable automata follows.

Consider a (symbolic) word $v = v_1 v_2 \dots v_n \in (\Sigma_B \cup \mathcal{X} \cup \{y\})^*$ recognized by B , and another word $w = w_1 w_2 \dots w_n \in \Sigma^*$. We say that w is a *legal instance* of v in $\mathcal{B}$ if the following holds:

- $v_i = w_i$ for every $v_i \in \Sigma_B$,
- For $v_i, v_j \in \mathcal{X}$, it holds that $w_i = w_j$ iff $v_i = v_j$, and $w_i, w_j \notin \Sigma_B$ and
- For $v_i = y$ and $v_j \neq y$, it holds that $w_i \neq w_j$.

Intuitively, a legal instance of v leaves all occurrences of $v_i \in \Sigma_B$ unchanged, associates every occurrence of $v_j \in \mathcal{X}$ with the same unique letter, not in Σ_B , and associates every occurrence of y freely with letters from $\Sigma \setminus \Sigma_B$, different from these associated with $\mathcal{X}$ variables.

Lemma 8. *For every variable automaton $\mathcal{B}$ we can construct a GVA $\mathcal{A}$ such that $L(\mathcal{A}) = L(\mathcal{B})$.*

Proof. Let $\mathcal{B} = \langle \Sigma, \mathcal{X} \cup \{y\}, B \rangle$ be a variable automaton with $B = \langle \Sigma_B \cup \mathcal{X} \cup \{y\}, Q, Q_0, \delta_B, F \rangle$ as pattern automaton. For a variable $x \in \mathcal{X}$, define the guards

$$\begin{aligned}\phi_x &= \bigwedge \{(x \neq z) \mid z \in \Sigma_B\} \\ \phi_y &= \bigwedge \{(y \neq z) \mid z \in \Sigma_B \cup \mathcal{X}\}\end{aligned}$$

Out of $\mathcal{B}$ we can construct an equivalent GVA $\mathcal{A} = \langle \Sigma, \mathcal{X} \cup \{y\}, Q, Q_0, \emptyset, \delta_{\mathcal{A}}, F, \kappa \rangle$, where the transition function $\delta_{\mathcal{A}}$ is defined by:

$$\begin{cases} q_2 \in \delta_{\mathcal{B}}(q_1, a) & \text{iff } q_2 \in \delta_{\mathcal{A}}(q_1, a), \text{ with } a \in \Sigma_B \\ q_2 \in \delta_{\mathcal{B}}(q_1, x) & \text{iff } q_2 \in \delta_{\mathcal{A}}(q_1, x, \phi_x), \text{ with } x \in \mathcal{X} \\ q_2 \in \delta_{\mathcal{B}}(q_1, y) & \text{iff } q_2 \in \delta_{\mathcal{A}}(q_1, y, \phi_y). \end{cases}$$

And the refreshing function κ is defined by:

$$\begin{cases} \kappa(x) = \emptyset, & \text{for all } x \in \mathcal{X}, \text{ and} \\ \kappa(y) = \{q \mid \exists q' \in Q, g \in \mathbb{G} \text{ s.t. } q' \in \delta_{\mathcal{A}}(q, y, g)\} \end{cases}$$

Claim 4 *Consider a word $v = v_1 v_2 \dots v_n \in (\Sigma_B \cup \mathcal{X} \cup \{y\})^*$ that reads along B , and another word $w = w_1 w_2 \dots w_n \in \Sigma^*$. There exists a run $q_0, M_0 \xrightarrow{\alpha_1, g_1} \dots \rightarrow q_n, M_n$ in $\mathcal{A}$ over w where $M_0 = \emptyset$ iff w is a legal instance of v in $\mathcal{B}$.*

Proof (Of Claim 4). The proof is by induction on n in both directions. The case $n = 0$ trivially holds. Assume that the claim holds up to n , and let's prove it for $n + 1$.

$\Rightarrow$) Assume a transition $q_{n+1} \in \delta_{\mathcal{A}}(q_n, \alpha_{n+1}, g_{n+1})$ in $\mathcal{A}$. From the definition of the transition function $\delta_{\mathcal{A}}$, we have $q_{n+1} \in \delta_{\mathcal{B}}(q_n, \alpha_{n+1})$. We distinguish three cases depending on α_{n+1} :

- If α_{n+1} is a letter in Σ , then this case is immediate.
- If α_{n+1} is a variable $x \in \mathcal{X}$, then in this case we recall that $g_{n+1} = \phi_x$, where ϕ_x is defined above. We distinguish two cases:
 - If $\forall i \leq n, \alpha_i \neq y$: in this case the transition in $\mathcal{B}$ is possible since ϕ_x implies that $w_{n+1} \notin \Sigma_B$.
 - Otherwise, that is, let $n_1, \dots, n_m \in \{1, \dots, n\}$ be all the indexes such that $\alpha_{n_l} = y$ for all $l = 1, \dots, m$. Since the variables x is never refreshed, then for $v_i, v_j \in \mathcal{X}$, it holds that $w_i = w_j$ iff $v_i = v_j$. From the definition of $\phi_{\alpha_n} = \bigwedge_{(b_k \in \Sigma_B)} (x \neq b_k)$, we get $w_i, w_j \notin \Sigma_B$. Thus w is a legal instance of v .
- If α_{n+1} is the variable y , then recall that y is refreshed in the state q_n of $\mathcal{A}$ and $g_{n+1} = \phi_y$, where ϕ_y is defined above. In this case we let $\mathcal{X} = \mathcal{X}' \uplus \text{dom}(M_n)$. Since $M_n \vdash \phi_y$, then for all $x \in \text{dom}(M_n)$ we have that $w_{n+1} \neq M_n(x)$, and for all $b_k \in \Sigma_B$, $w_{n+1} \neq b_k$. Hence for all $v_j \neq y$ we have that $w_i \neq w_j$.

$\Leftarrow$) Assume a transition $q_{n+1} \in \delta_{\mathcal{B}}(q_n, \alpha_{n+1})$. From the definition of the transition function $\delta_{\mathcal{A}}$, we have that $q_{n+1} \in \delta_{\mathcal{A}}(q_n, \alpha_{n+1}, g_{n+1})$ in $\mathcal{A}$, for some guard $g_{n+1} \in \mathbb{G}$. We distinguish three cases depending on α_{n+1} :

- If α_{n+1} is a letter in Σ , then in this case $g_{n+1} = \text{true}$, and we are done.
- If α_{n+1} is a variable $x \in \mathcal{X}$, then in this case we have that for $v_i, v_j \in \mathcal{X}$, it holds that $w_i = w_j$ iff $v_i = v_j$, and $w_i, w_j \notin \Sigma_B$. We recall that $g_{n+1} = \phi_x$, where ϕ_x is defined above. We distinguish two cases:

- If $\forall i < n, \alpha_i \neq y$: in this case the transition in $\mathcal{A}$ is possible since $w_{n+1} \notin \Sigma_B$ implies that ϕ_x .
- Otherwise, that is, let $n_1, \dots, n_m \in \{1, \dots, n\}$ be all the indexes such that $\alpha_{n_j} = y$. In this case Since for all $v_i, v_j \in \mathcal{X}$, it holds that $w_i = w_j$ iff $v_i = v_j$, and $w_i, w_j \notin \Sigma_B$ then ϕ_x holds as well.
- If α_n is the variable y , then for all $v_j \neq y$, it holds that $w_i \neq w_j$. Since $g_{n+1} = \phi_y$, then $M_n \vdash \phi_y$. Thus, the transition in $\mathcal{A}$ is possible.

□

□

D.2 GVAs vs. FMA

We recall the formal definition of the computations of FMA.

Definition 14. Let $\mathcal{F} = \langle \Sigma, Q, q_0, \tau, \varrho, \delta, F \rangle$ be a FMA. A configuration is a pair (q, R) where $q \in Q$ and $\{1, \dots, k\} : \mathcal{X} \rightarrow \Sigma$ is an assignment. We define a transition relation over the configurations as follows: let $a \in \Sigma$, then, $(q, R) \xrightarrow{a} (q', R')$ if there exists $i \in \{1, \dots, k\}$ and $q' \in \delta(q, i)$ such that:

1. $a = R(i)$ and $R = R'$, or
2. $a \notin \text{codom}(R)$, $\varrho(q) = i$, $R'(i) = a$, and $R(j) = R'(j)$ for all $j \neq i$.

A finite word $w = w_1 w_2 \dots w_n \in \Sigma^*$ is recognized by $\mathcal{F}$ iff there exists a run $(q_0, R_0) \xrightarrow{w_1} (q_1, R_1) \xrightarrow{w_2} \dots \xrightarrow{w_n} (q_n, R_n)$, such that $R_0 = \tau$, $q_0 \in Q_0$ and $q_n \in F$.

Before proving that GVAs are more expressive than FMA (i.e. Lemma 9 below), let's give the idea behind the proof: We next show that for every FMA with k registers and n states, we can construct a GVA with $k + 1$ variables and $O(n \cdot (k + 1)!)$ states recognizing the same language. The idea is that k variables, say $\{x_1, \dots, x_k\}$, are used to mimic the k registers, say $\{1, \dots, k\}$, and the remaining variable, say x_{k+1} , is used to store the current letter.

The problem is, while a FMA checks that the current letter does not occur in any register and rewrites this letter in a register (i.e. Item 2 in Def. 14), a GVA cannot—a priori—mimic these two operations through only one transition. Our solution consists changing the correspondence between the k registers and the $k + 1$ variables. That is, instead of copying the value of the extra variable x_{k+1} (i.e. the current letter) to the variable x_i that corresponds to the register i , the role of the variables x_i and x_{k+1} is swapped: the variable x_{k+1} will correspond to the register i , and

x_i will play the role of the extra variable that will be used for storing the current letter.

For the formal proof, we need to introduce some notations. Let $k \in \mathbb{N}$. Let $\mathcal{X} = \{x_1, \dots, x_{k+1}\}$. We denote by $\widehat{\Psi}^{k, \mathcal{X}}$ the set of all injective total functions from $[k]$ to $\mathcal{X}$. We define $\widehat{\psi}$ by $\widehat{\psi}(i) = x_i$ for all $i \in [k]$. If $\psi \in \Psi^{k, \mathcal{X}}$, then we denote by $\mathcal{X}(\psi)$ the unique variable x such that there is no $j \in [k]$ where $\psi(j) = x$. Thus, for a fixed $i \in [k]$, we define the function $\psi^i : \mathbb{N} \rightarrow \mathcal{X}$ by $\psi^i(i) = \mathcal{X}(\psi)$ and $\psi^i(j) = \psi(j)$ for all $j \neq i$. If $M : \mathcal{X} \rightarrow \Sigma$ is a substitution, $R : [k] \rightarrow \Sigma$ is an assignment, and $\psi \in \Psi^{k, \mathcal{X}}$, then we shall write $M \cong_\psi R$ iff $|\text{dom}(M)| = |\text{dom}(R)|$ and for all $j \in [k]$ we have that $M(\psi(j)) = R(j)$.

Lemma 9. *For every FMA $\mathcal{F}$ with k registers, there exists a GVA $\mathcal{F}'$ with $k+1$ variables such that $L(\mathcal{F}') = L(\mathcal{F})$.*

Proof. Let $\mathcal{F} = \langle \Sigma, k, Q, q_0, \tau, \varrho, \delta, F \rangle$ be a FMA. We define $\mathcal{F}'$ to be the GVA $\mathcal{F}' = \langle \Sigma, \mathcal{X}, Q', q'_0, \tau', \delta', F', \kappa \rangle$ defined by:

$$\begin{cases} \mathcal{X} = \{x_1, \dots, x_{k+1}\} \\ Q' = \{(q, \psi) \mid q \in Q \text{ and } \psi \in \widehat{\Psi}^{k, \mathcal{X}}\} \\ q'_0 = (q_0, \widehat{\psi}) \\ \tau' = \{\widehat{\psi}(j) \mapsto a_j \mid \tau(j) = a_j \ \forall j \in [k]\} \\ F' = \{(q, \psi) \mid q \in F \text{ and } \psi \in \Psi^{k, \mathcal{X}}\} \end{cases}$$

and

$$\begin{cases} (q', \psi) & \in \delta'((q, \psi), \mathcal{X}(\psi), (\mathcal{X}(\psi) = \psi(i))) \text{ and} \\ (q', \psi^i) & \in \delta'((q, \psi), \mathcal{X}(\psi), \bigwedge_{j \in [k]} (\mathcal{X}(\psi) \neq \psi(j))) \text{ with } i = \varrho(q) \\ \text{iff} & \\ q' & \in \delta(q, i) \end{cases} \quad (10)$$

Finally, the refreshing function κ is defined by $\kappa(x) = \{(q, \psi) \mid x = \mathcal{X}(\psi)\}$.

Fact 5 *Let $\mathbf{w} = w_1 \dots w_n$ be a word in Σ^n . There exists a run*

$$(q_0, \psi_0), M_0 \xrightarrow{\alpha_1, g_1} \dots \xrightarrow{\alpha_n, g_n} (q_n, \psi_n), M_n$$

in $\mathcal{F}'$ over $\mathbf{w}$, in which $M_0 = \varrho'$ and $\psi_0 = \widehat{\psi}$, iff there is a run

$$q_0, R_0 \xrightarrow{w_1} \dots \xrightarrow{w_n} q_n, R_n$$

in $\mathcal{F}$ over $\mathbf{w}$, in which $R_0 = \varrho$, such that $M_i \cong_{\psi_i} R_i$ for all $i = 0, \dots, n$.

Proof (Of Fact 5). By induction on n . The case $n = 0$ follows from the definition of ϱ' and $\widehat{\psi}$. In other words, we have that $\varrho' \cong_{\widehat{\psi}} \varrho$. Assume the claim holds up to n . Let us prove the equivalence for $n + 1$.

$\Rightarrow$) Assume $(q_{n+1}, \psi_{n+1}) \in \delta'((q_n, \psi_n), \mathcal{X}(\psi_n), g_n)$. We distinguish two cases depending on the nature of this transition, i.e. depending on g_n :

- If $g_n = (\mathcal{X}(\psi_n) = \psi(i))$ then it follows from the definition of δ' in Eq (10) that $\psi_{n+1} = \psi_n$ and $(q_n, \psi_n) \in \kappa(\mathcal{X}(\psi_n))$ and $q_{n+1} \in \delta(q_n, i)$. On the one hand, we have that $M_{n+1} = M_n$ since the variable $\mathcal{X}(\psi_n)$ is refreshed in both states (q_n, ψ_n) and (q_{n+1}, ψ_n) . From the induction hypothesis we have $M_n \cong_{\psi_n} R_n$. Thus $M(\psi(i)) = R(i) = w_{n+1}$. It follows from the definition of the configurations for FMA that the Case 1 in Def. 14 is applicable and hence $R_n = R_{n+1}$. Finally, the invariant $M_{n+1} \cong_{\psi_{n+1}} R_{n+1}$ holds, since $M_n = M_{n+1}$ and $R_{n+1} = R_n$ and $\psi_{n+1} = \psi_n$.
- If $g_n = \bigwedge_{j \in [k]} (\psi_n(j) \neq \mathcal{X}(\psi_n))$ then it follows from the definition of δ' in Eq (10) that $\psi_{n+1} = \psi_n^i$ and $q_{n+1} \in \delta(q_n, i)$ and $\varrho(p_n) = i$. From the induction hypothesis we have that $M_n \cong_{\psi_n} R_n$. Since $M_n \uplus \{\mathcal{X}(\psi_n) \mapsto w_{n+1}\} \models \bigwedge_{j \in [k]} (x_j \neq \mathcal{X}(\psi_n))$ and $\mathcal{X}(\psi_n) \notin \text{dom}(M_n)$ then for all $j \in [k]$ we have $M_n(\psi_n(j)) \neq w_{n+1}$. Hence,

$$R(j) \neq R(s) \quad \forall j \neq s, \text{ and } R_n(i) = M_n(x_{\psi_n(i)}) \quad (11)$$

It follows from the definition of the configurations for FMA that the Case 2 in Def. 14 is applicable. Hence $R_{n+1}(j) = R_n(j)$ for all $j \neq i$ and $R_{n+1}(i) = w_{n+1}$. It remains to show that $M_{n+1} \cong_{\psi_{n+1}} R_{n+1}$, where $\psi_{n+1} \stackrel{\text{def}}{=} \psi_n^i$.

On the one hand, from the definition of the run for GVAs we have

$$\begin{cases} M_{n+1}(\psi_n(i)) &= w_{n+1}, \text{ and} \\ M_{n+1}(\psi_n(j)) &= M_n(\psi_n(j)) \quad \forall j \in [k] \setminus \{i\} \end{cases} \quad (12)$$

On the other hand, it follows from the definition of ψ_n^i that

$$\begin{cases} \psi_n^i(i) &= \mathcal{X}(\psi_n), \text{ and} \\ \psi_n^i(j) &= \psi_n(j) \quad \forall j \in [k] \setminus \{i\} \end{cases} \quad (13)$$

From (12) and (13) we get

$$\begin{cases} M_{n+1}(\psi_{n+1}(i)) &= M_{n+1}(\psi_n^i(i)) = M_{n+1}(\mathcal{X}(\psi_n)) = w_{n+1} = R_{n+1}(i), \text{ and} \\ M_{n+1}(\psi_{n+1}(j)) &= M_{n+1}(\psi_n^i(j)) = M_{n+1}(\psi_n(j)) \\ &= M_n(\psi_n(j)) = R_{n+1}(j) \quad \forall j \in [k] \setminus \{i\} \end{cases}$$

Hence, $M_{n+1} \cong_{\psi_{n+1}} R_{n+1}$.

$\Leftarrow$) Assume $q_{n+1} \in \delta(q_n, i)$. We distinguish two cases depending on the transition made in $\mathcal{F}$ (i.e. the same two cases in Def. 14):

- If $w_{n+1} = R_n(i)$ and in this case $R_{n+1} = R_n$. From the definition of δ' in Eq (10) we have $(q_{n+1}, \psi_{n+1}) \in \delta'((q_n, \psi_n), \mathcal{X}(\psi_n), (\mathcal{X}(\psi_n) = \psi_n(i)))$. Since the variable $\mathcal{X}(\psi_n)$ is free in state (q_n, ψ_n) then the latter transition is possible. Thus we get $M_{n+1} = M_n$ and $\psi_{n+1} = \psi_n$. From the induction hypothesis we have $M_n \cong_{\psi_n} R_n$. Hence, $M_{n+1} \cong_{\psi_{n+1}} R_{n+1}$.
- If $w_{n+1} \notin \text{codom}(R_n)$, $\varrho(p) = i$, and in this case $R_n(j) = R_{n+1}(j)$ for all $j \neq i$. From the definition of δ' in Eq (10) we have

$$(q_{n+1}, \psi_{n+1}^i) \in \delta'((q_n, \psi_n), \mathcal{X}(\psi_n), \bigwedge_{j \in [k]} (\mathcal{X}(\psi_n) \neq \psi_n(j)))$$

We show that this transition is possible. Since the variable $\mathcal{X}(\psi_n)$ is refreshed in the state (q_n, ψ_n) , then we must show that $M_n \models \bigwedge_{j \in [k]} (w_{n+1} \neq \psi_n(j))$, i.e. that $\bigwedge_{j \in [k]} (w_{n+1} \neq M_n(\psi_n(j)))$ holds. But since $w_{n+1} \notin \text{codom}(R_n)$ and $M_n \cong_{\psi_n} R_n$, then $w_{n+1} \notin \text{codom}(M_n)$ as well. Thus, $\bigwedge_{j \in [k]} (w_{n+1} \neq M_n(\psi_n(j)))$ holds. It remains to show that $M_{n+1} \cong_{\psi_{n+1}} R_{n+1}$. But this is the same as the proof of the second case of the direction ($\Rightarrow$).

□

□